### Attachment A. Proposed Rule Amendments Draft Rules – Edits Highlighted

**Key to Identifying Changed Text:** 

Deleted Text New/inserted text

### Division 41 WATER QUALITY STANDARDS: BENEFICIAL USES, POLICIES, AND CRITERIA FOR OREGON

### 340-041-0033 Toxic Substances

- (1) Toxic Substances Narrative. Toxic substances may not be introduced above natural background levels in waters of the state in amounts, concentrations, or combinations that may be harmful, may chemically change to harmful forms in the environment, or may accumulate in sediments or bioaccumulate in aquatic life or wildlife to levels that adversely affect public health, safety, or welfare or aquatic life, wildlife or other designated beneficial uses.
- (2) Aquatic Life Numeric Criteria. Levels of toxic substances in waters of the state may not exceed the applicable aquatic life criteria as defined in Table 30 under OAR 340-041-8033.
- (3) Human Health Numeric Criteria. The criteria for waters of the state listed in Table 40 under OAR 340-041-8033 are established to protect Oregonians from potential adverse health effects associated with long-term exposure to toxic substances associated with consumption of fish, shellfish and water.
- (4) To establish permit or other regulatory limits for toxic substances without criteria in Table 30 under OAR 340-041-8033 or Table 40 under 340-041-8033, DEQ may use the guidance values in Table 31 under 340-041-8033, public health advisories, and published scientific literature. DEQ may also require or conduct bio-assessment studies to monitor the toxicity to aquatic life of complex effluents, other suspected discharges or chemical substances without numeric criteria.
- (5) Establishing Site-Specific Background Pollutant Criteria: This provision is a performance- based water quality standard that results in site-specific human health water quality criteria under the conditions and procedures specified in this rule section. It addresses existing permitted discharges of a pollutant removed from the same body of water. For waterbodies where a discharge does not increase the pollutant's mass and

does not increase the pollutant concentration by more than 3 percent, and where the water body meets a pollutant concentration associated with a risk level of 1 x 10-4, DEQ concludes that the pollutant concentration continues to protect human health.

- (a) Definitions: As used in this section:
- (A) "Background pollutant concentration" means the ambient water body concentration immediately upstream of the discharge, regardless of whether those pollutants are natural or result from upstream human activity.
- (B) An "intake pollutant" is the amount of a pollutant present in waters of the state (including groundwater) as provided in subsection (C), below, at the time it is withdrawn from such waters by the discharger or other facility supplying the discharger with intake water.
- (C) "Same body of water": An intake pollutant is considered to be from the "same body of water" as the discharge if DEQ finds that the intake pollutant would have reached the vicinity of the outfall point in the receiving water within a reasonable period had the permittee not removed it. To make this finding, DEQ requires information showing that:
- (i) The background concentration of the pollutant in the receiving water (excluding any amount of the pollutant in the facility's discharge) is similar to that in the intake water; and,
- (ii) There is a direct hydrological connection between the intake and discharge points.
- (I) DEQ may also consider other site-specific factors relevant to the transport and fate of the pollutant to make the finding in a particular case that a pollutant would or would not have reached the vicinity of the outfall point in the receiving water within a reasonable period had the permittee not removed it.
- (II) An intake pollutant from groundwater may be considered to be from the "same body of water" if DEQ determines that the pollutant would have reached the vicinity of the outfall point in the receiving water within a reasonable period had the permittee not removed it. A pollutant is not from the same body of water if the groundwater contains the pollutant partially or entirely due to past or present human activity, such as industrial, commercial, or municipal operations, disposal actions, or treatment processes.
- (iii) Water quality characteristics (e.g., temperature, pH, hardness) are similar in the intake and receiving waters.
- (b) Applicability
- (A) DEQ may establish site-specific criteria under this rule section only for carcinogenic pollutants.

- (B) Site-specific criteria established under this rule section apply in the vicinity of the discharge for purposes of establishing permit limits for the specified permittee.
- (C) The underlying waterbody criteria continue to apply for all other Clean Water Act programs.
- (D) The site-specific background pollutant criterion will be effective when DEQ issues the permit for the specified permittee.
- (E) DEQ will reevaluate any site-specific criteria developed under this procedure upon permit renewal.
- (c) DEQ may establish a site-specific background pollutant criterion when all of the following conditions are met:
- (A) The discharger has a currently effective NPDES permit;
- (B) The mass of the pollutant discharged to the receiving waterbody does not exceed the mass of the intake pollutant from the same body of water, as defined in section (5)(a)(C) above, and therefore does not increase the total mass load of the pollutant in the receiving water body;
- (C) DEQ has not assigned the discharger a TMDL wasteload allocation for the pollutant in question;
- (D) The permittee uses any feasible pollutant reduction measures available and known to minimize the pollutant concentration in their discharge;
- (E) The pollutant discharge has not been chemically or physically altered in a manner that causes adverse water quality impacts that would not occur if the intake pollutants were left in-stream; and,
- (F) The timing and location of the pollutant discharge would not cause adverse water quality impacts that would not occur if the intake pollutant were left in-stream.
- (d) The site-specific background pollutant criterion must be the most conservative of the following four values. Section (5)(e) of this rule describes the procedures for deriving these values.
- (A) The projected in-stream pollutant concentration resulting from the current discharge concentration and any feasible pollutant reduction measures under (c)(D) above, after mixing with the receiving stream.
- (B) The projected in-stream pollutant concentration resulting from the portion of the current discharge concentration associated with the intake pollutant mass after mixing with the receiving stream. This analysis ensures that there will be no increase in the

mass of the intake pollutant in the receiving water body as required by condition (c)(B) above.

- (C) The projected in-stream pollutant concentration associated with a 3 percent increase above the background pollutant concentration as calculated:
- (i) For the main stem Willamette and Columbia Rivers, using 25 percent of the harmonic mean flow of the waterbody.
- (ii) For all other waters, using 100 percent of the harmonic mean flow or similar critical flow value of the waterbody.
- (D) A criterion concentration value representing a human health risk level of 1 x 10-4. DEQ calculates this value using EPA's human health criteria derivation equation for carcinogens (EPA 2000), a risk level of 1 x 10-4, and the same values for the remaining calculation variables that were used to derive the underlying human health criterion.
- (e) Procedure to derive a site-specific human health water quality criterion to address a background pollutant:
- (A) DEQ will develop a flow-weighted characterization of the relevant flows and pollutant concentrations of the receiving waterbody, effluent and all facility intake pollutant sources to determine the fate and transport of the pollutant mass.
- (i) The pollutant mass in the effluent discharged to a receiving waterbody may not exceed the mass of the intake pollutant from the same body of water.
- (ii) Where a facility discharges intake pollutants from multiple sources that originate from the receiving waterbody and from other waterbodies, DEQ will calculate the flow-weighted amount of each source of the pollutant in the characterization.
- (iii) Where a municipal water supply system provides intake water for a facility and the supplier provides treatment of the raw water that removes an intake water pollutant, the concentration and mass of the intake water pollutant must be determined at the point where the water enters the water supplier's distribution system.
- (B) Using the flow weighted characterization developed in section (5)(e)(A), DEQ will calculate the in-stream pollutant concentration following mixing of the discharge into the receiving water. DEQ will use the resultant concentration to determine the conditions in section (5)(d)(A) and (B).
- (C) Using the flow-weighted characterization, DEQ will calculate the in-stream pollutant concentration based on an increase of 3 percent above background pollutant concentration. DEQ will use the resultant concentration to determine the condition in Section (5)(d)(C).

- (i) For the main stem Willamette and Columbia Rivers, DEQ will use 25 percent of the harmonic mean flow of the waterbody.
- (ii) For all other waters, DEQ will use 100 percent of the harmonic mean flow or similar critical flow value of the waterbody.
- (D) DEQ will select the most conservative of the following values as the site-specific water quality criterion.
- (i) The projected in-stream pollutant concentration described in section (5)(e)(B);
- (ii) The in-stream pollutant concentration based on an increase of 3 percent above background described in section (5)(e)(C); or
- (iii) A water quality criterion based on a risk level of 1 x 10-4.
- (f) Calculation of water quality based effluent limits based on a site-specific background pollutant criterion:
- (A) For discharges to receiving waters with a site-specific background pollutant criterion, DEQ will use the site-specific criterion in the calculation of a numeric water quality based effluent limit.
- (B) DEQ will compare the calculated water quality based effluent limits to any applicable aquatic toxicity or technology based effluent limits and select the most conservative for inclusion in the permit conditions.
- (g) In addition to the water quality based effluent limits described in section (5)(f), DEQ will calculate a mass-based limit where necessary to ensure that the condition described in section (5)(c)(B) is met. Where mass-based limits are included, the permit will specify how DEQ will assess compliance with mass-based effluent limitations.
- (h) The permit shall include a provision requiring DEQ to consider the re-opening of the permit and re-evaluation of the site-specific background pollutant criterion if new information shows the discharger no longer meets the conditions described in subsections (5)(c) and (e).
- (i) Public Notification Requirements.
- (A) If DEQ proposes to grant a site-specific background pollutant criterion, it must provide public notice of the proposal and hold a public hearing. The public notice may be included in the public notification of a draft NPDES permit or other draft regulatory decision that would rely on the criterion and will also be published on DEQ's water quality standards website;

- (B) DEQ will publish a list of all site-specific background pollutant criteria approved according to this rule. DEQ will add the criterion to this list within 30 days of its effective date. The list will identify the:
- (i) Permittee;
- (ii) Site-specific background pollutant criterion and the associated risk level;
- (iii) Waterbody to which the criterion applies;
- (iv) Allowable pollutant effluent limit; and,
- (v) How to obtain additional information about the criterion.
- (6) Arsenic Reduction Policy: The inorganic arsenic criterion for the protection of human health from the combined consumption of organisms and drinking water is 2.1 micrograms per liter. While this criterion is protective of human health and more stringent than the federal maximum contaminant level (MCL) for arsenic in drinking water, which is 10 micrograms per liter, it is based on a higher risk level than EQC used to establish other human health criteria. This higher risk level recognizes that much of the risk is due to naturally high levels of inorganic arsenic in Oregon's waterbodies. In order to maintain the lowest human health risk from inorganic arsenic in drinking water, EQC determined that it is appropriate to adopt the following policy to limit the human contribution to that risk.
- (a) It is EQC policy to reduce the addition of inorganic arsenic from new or existing anthropogenic sources to waters of the state within a surface water drinking water protection area to the maximum amount feasible. The requirements of this rule section (OAR 340-041-0033(6)) apply to sources that discharge to surface waters of the state with an ambient inorganic arsenic concentration equal to or lower than the applicable numeric inorganic arsenic criteria for the protection of human health.
- (b) Definitions. As used in this section:
- (A) "Add inorganic arsenic" means to discharge a net mass of inorganic arsenic from a point source (the mass of inorganic arsenic discharged minus the mass of inorganic arsenic taken into the facility from a surface water source).
- (B) A "surface water drinking water protection area," means an area delineated as such by DEQ under the source water assessment program of the federal Safe Drinking Water Act, 42 U.S.C. § 300j 13. DEQ delineates these areas to protect public or community drinking water supplies that use surface water sources. These delineations are on DEQ's drinking water program Web page.
- (C) "Potential to significantly increase inorganic arsenic concentrations in the public drinking water supply source water" means:

- (i) A discharge will increase the concentration of inorganic arsenic in the receiving water by 10 percent or more after mixing with the harmonic mean flow of the receiving water; or
- (ii) As an alternative, if sufficient data are available, the discharge will increase the concentration of inorganic arsenic in the surface water intake water of a public water system by 0.021 micrograms per liter or more based on a mass balance calculation.
- (c) Following the effective date of this rule, applications for an individual NPDES permit or permit renewal received from industrial dischargers located in a surface water drinking water protection area and identified by DEQ as likely to add inorganic arsenic to the receiving water must include sufficient data to enable DEQ to determine whether:
- (A) The discharge adds inorganic arsenic; and,
- (B) The discharge has the potential to significantly increase inorganic arsenic concentrations in the public drinking water supply source water.
- (d) Where DEQ determines that both conditions in subsection (c) of this section (6) are true, the industrial discharger must develop an inorganic arsenic reduction plan and propose all feasible measures to reduce its inorganic arsenic loading to the receiving water. The proposed plan, including proposed measures, monitoring and reporting requirements, and a schedule for those actions, will be described in the fact sheet and incorporated into the source's NPDES permit after public comment and DEQ review and approval. In developing the plan, the source must:
- (A) Identify how much it can minimize its inorganic arsenic discharge through pollution prevention measures, process changes, wastewater treatment, alternative water supply for groundwater users, or other possible pollution prevention and control measures;
- (B) Evaluate the costs, feasibility and environmental impacts of the potential inorganic arsenic reduction and control measures;
- (C) Estimate the predicted reduction in inorganic arsenic and the reduced human health risk expected to result from the control measures;
- (D) Propose specific inorganic arsenic reduction or control measures, if feasible, and an implementation schedule; and,
- (E) Propose monitoring and reporting requirements to document progress in plan implementation and the inorganic arsenic load reductions.
- (e) In order to implement this section, DEQ will develop the following information and guidance within 120 days of the effective date of this rule and periodically update it as warranted by new information:

- (A) A list of industrial sources or source categories, including industrial stormwater and sources covered by general permits likely to add inorganic arsenic to surface waters of the state. For industrial sources or source categories permitted under a general permit that have been identified by DEQ as likely sources of inorganic arsenic, DEQ will evaluate options for reducing inorganic arsenic during permit renewal or evaluation of Stormwater Pollution Control Plans.
- (B) Quantitation limits for monitoring inorganic arsenic concentrations.
- (C) Information and guidance to assist sources in estimating, according to subsection (d)(C) of this section, the reduced human health risk expected to result from inorganic arsenic control measures based on the most current EPA risk assessment.
- (f) It is the policy of EQC that landowners engaged in agricultural or development practices on land where pesticides, fertilizers, or soil amendments containing arsenic are currently being or have previously been applied, implement conservation practices to minimize the erosion and runoff of inorganic arsenic to waters of the state or to a location where such material could readily migrate into waters of the state.

NOTE: Tables 30, 31 and 40 are found under OAR 340-041-8033.

**Statutory/Other Authority:** ORS 468.020, 468B.030, 468B.035 & 468B.048 **Statutes/Other Implemented:** ORS 468B.030, 468B.035 & 468B.048 **History:** 

DEQ 11-2016, f. & cert. ef. 11-2-16

DEQ 1-2015, f. & cert. ef. 1-7-15

DEQ 17-2013, f. 12-23-13, cert. ef. 4-18-14

DEQ 10-2011, f. & cert. ef. 7-13-11

DEQ 8-2011, f. & cert. ef. 6-30-11

DEQ 17-2010, f. & cert. ef. 12-21-10

DEQ 3-2004, f. & cert. ef. 5-28-04

DEQ 17-2003, f. & cert. ef. 12-9-03

### Division 41 WATER QUALITY STANDARDS: BENEFICIAL USES, POLICIES, AND CRITERIA FOR OREGON

### 340-041-8033 Division 41 Tables and Figures

- (1) Table 30: Aquatic Life Water Quality Criteria for Toxic Pollutants. This table, referenced in OAR 340-041-0033, contains information about the applicability and content of the criteria contained in the table.
- (2) Table 31: Aquatic Life Water Quality Guidance Values for Toxic Pollutants. This table, referenced in OAR 340-041-0033, contains information about the applicability and content of the criteria contained in the table.
- (32) Table 40: Human Health Water Quality Criteria for Toxic Pollutants. This table, referenced in OAR 340-041-0033, contains information about the applicability and content of the criteria contained in the table.

[ED. NOTE: To view attachments referenced in rule text, click here for PDF copy.] **Statutory/Other Authority**: ORS 468.020, 468B.030, 468B.035 & 468B.048Statutes/Other **Implemented**: ORS 468B.030, 468B.035 & 468B.048 **History**:

DEQ 13-2019, amend filed 05/16/2019, effective 05/16/2019

DEQ 13-2017, minor correction filed 10/30/2017, effective 10/30/2017

DEQ 11-2016, f. & cert. ef. 11-2-16

DEQ 1-2015, f. & cert. ef. 1-7-15



The concentration for each compound listed in Table 30 is a criterion established for waters of the state in order to protect aquatic life. The aquatic life criteria apply to waterbodies where the protection of fish and aquatic life is a designated use. All values are expressed as micrograms per liter ( $\mu$ g/L). Compounds are listed in alphabetical order with the corresponding information: the Chemical Abstract Service (CAS) number, whether there is a human health criterion for the pollutant (i.e. "y"= yes, "n" = no), and the associated aquatic life freshwater and saltwater acute and chronic criteria. *Italicized* pollutants are not identified as priority pollutants by EPA. Dashes in the table column indicate that there is no aquatic life criterion.

Unless otherwise noted in the table below, the acute criterion is the Criterion Maximum Concentration (CMC) applied as a one-hour average concentration, and the chronic criterion is the Criterion Continuous Concentration (CCC) applied as a 96-hour (4 days) average concentration. The CMC and CCC criteria may not be exceeded more than once every three years. Footnote A, associated with eleven pesticide pollutants in Table 30, describes the exception to the frequency and duration of the toxics criteria stated in this paragraph.

	Aquatic Elic Water Quality Official for Toxic Foliatants										
				Freshwater (µg/L)		Saltwater (µg/L)					
No.	Pollutant	CAS Number	Human Health Criterion	Acute Criterion (CMC)	Chronic Criterion (CCC)	Acute Criterion (CMC)	Chronic Criterion (CCC)				
1	<u>Acrolein</u>	107028	У	3.0	3.0	Ш	Н				
NOT	E: These acrolein c	riteria are n	ot applicable	e for Clean Wat	ter Act purpose	es until approve	ed by EPA.				
<u>2</u> 1	Aldrin	309002	у	3 A		1.3 <sup>A</sup>					
<sup>A</sup> Sec	<sup>A</sup> See expanded endnote A at bottom of Table 30 for alternate frequency and duration of this criterion.										
<u>3</u> 2	Alkalinity		n		20,000 B						

				Fresh (µg			twater <i>ıg/L)</i>	
No.	Pollutant	CAS Number	Human Health Criterion	Acute Criterion (CMC)	Chronic Criterion (CCC)	Acute Criterion (CMC)	Chronic Criterion (CCC)	
B Criterion shown is the minimum (i.e. CCC in water may not be below this value in order to protect aquatic life).								
4	Aluminum	7429905	n	See O, P	See O, P			

O The freshwater criterion for aluminum is a function of the pH, dissolved organic carbon, and total hardness in the water column. Acute (CMC) and chronic (CCC) freshwater aluminum criteria values shall be calculated using the 2018 Aluminum Criteria Calculator (Aluminum Criteria Calculator V.2.0.xlsx), or a calculator in R or other software package using the same 1985 Guidelines calculation approach and underlying model equations as in the Aluminum Criteria Calculator V.2.0.xlsx, as defined in EPA's Final Aquatic Life Ambient Water Quality Criteria for Aluminum (EPA 822-R-18-001). See also endnote O for procedures and information.

P Oregon will use analytical methods that measure the bioavailable fraction of aluminum unless total recoverable aluminum measurements are required by Federal regulations.

Note: These aluminum criteria are not applicable for CWA purposes until approved by EPA and, if necessary, EPA withdraws the federal criteria for Oregon. The currently applicable federally promulgated criteria may be found at 40 CFR 131.47.

<u>5</u> 3	Ammonia	7664417	n	The ammonia criteria are pH and temperature dependent — See ammonia criteria Tables 30(a)-(c) at end of Table 30. <sup>M</sup>	The ammonia criteria are pH, temperature and salinity dependent. Values for saltwater criteria (total ammonia) can be calculated from the tables specified in Ambient Water Quality Criteria for Ammonia (Saltwater)—1989 (EPA 440/5-88-004) See DEQ's calculator for calculating saltwater
					ammonia criteria at:

				Fresh (µg		Saltwater (µg/L)		
No.	Pollutant	CAS Number	Human Health Criterion	Acute Criterion (CMC)	Chronic Criterion (CCC)	Acute Criterion (CMC)	Chronic Criterion (CCC)	
				·		http://www.deq.state.or.us/w q/standards/toxics.htm		

M The acute criteria in Table 30(a) apply in waterbodies where salmonids are a designated use in OAR 340-041-0101 through OAR 340-041-0340. The acute criteria in Table 30(b) apply in waterbodies where salmonids are not a designated use. The chronic criteria in Table 30(c) apply where fish and aquatic life is a designated use. It is not necessary to account for the presence or absence of salmonids or the presence of any early life stage of fish for the chronic criteria. Refer to DEQ's beneficial use website at: http://www.deq.state.or.us/wq/standards/uses.htm for additional information on salmonid beneficial use designations, including tables and maps.

<u>6</u> 4	Arsenic	7440382	у	340 <sup>c, d</sup>	150 <sup>C, D</sup>	69 <sup>c, d</sup>	36 C, D		
<sup>c</sup> Criterion is expressed in terms of "dissolved" concentrations in the water column. <sup>D</sup> Criterion is applied as total inorganic arsenic (i.e. arsenic (III) + arsenic (V)).									
<u>7</u> 5	BHC Gamma (Lindane)	58899	у	0.95	0.08 <sup>A</sup>	0.16 <sup>A</sup>			
A See expanded endnote A at bottom of Table 30 for alternate frequency and duration of this criterion.									
<u>8</u> 6	Cadmium	7440439	n	See <mark>E₋C, F</mark>	See C, F	40- <u>33</u> c	8.8 <u>7.9</u> c		

<sup>&</sup>lt;sup>c</sup> Criterion is expressed in terms of "dissolved" concentrations in the water column.

NOTE: The federally promulgated freshwater acute cadmium criterion in 40 CFR 131.46 remains applicable for CWA purposes until EPA approves the freshwater acute cadmium criterion specified in Table 30 Endnotes C and F. The following saltwater cadmium criteria remain applicable until EPA approves the revised criteria in Table 30: the cadmium acute (CMC) is 40 µg/L and the chronic (CCC) is 8.8 µg/L, both expressed in terms of "dissolved" concentrations in the water column.

E The freshwater criterion for this metal is expressed as "total recoverable" and is a function of hardness (mg/L) in the water column. To calculate the criterion, use formula under expanded endnote E at bottom of Table 30.

F The freshwater criterion for this metal is expressed as a function of hardness (mg/L) in the water column. To calculate the criterion, use the formula under expanded endnote F at bottom of Table 30.

					Freshwater (μg/L)		Saltwater (µg/L)	
No.	Pollutant	CAS Number	Human Health Criterion	Acute Criterion (CMC)	Chronic Criterion (CCC)	Acute Criterion (CMC)	Chronic Criterion (CCC)	
9	<u>Carbaryl</u>	63252	<u>n</u>	<u>2.1</u>	<u>2.1</u>	<u>1.6</u>	==	
NOTE: These carbaryl criteria are not applicable for Clean Water Act purposes until approved by EPA.								
<u>10</u> 7	Chlordane	57749	у	2.4 <sup>A</sup>	0.0043 A	0.09 A	0.004 <sup>A</sup>	
<sup>A</sup> See	e expanded endnot	e A at botto	m of Table 3	30 for alternate	frequency and	duration of this	s criterion.	
<u>11</u> 8	Chloride	1688700 6	n	860,000	230,000			
<u>12</u> 9	Chlorine	7782505	n	19	11	13	7.5	
134 0	Chlorpyrifos	2921882	n	0.083	0.041	0.011	0.0056	
<u>14</u> 1 1	Chromium III	1606583 1	n	See C, F	See C, F			
F Th	erion is expressed e freshwater criterionalculate the criterion	on for this m	etal is expre	essed as a func	tion of hardnes	ss (mg/L) in the	e water column.	
154 2	Chromium VI	1854029 9	n	16 <sup>c</sup>	11 <sup>c</sup>	1100 <sup>c</sup>	<sub>50</sub> c	
<sup>c</sup> Crit	erion is expressed	in terms of '	'dissolved" c	oncentrations i	n the water co	lumn.		
<u>16</u> 4 3	Copper	7440508	у	See C, N	See C, N	4.8 <sup>c</sup>	3.1 <sup>C</sup>	

<sup>&</sup>lt;sup>c</sup> Criterion is expressed in terms of "dissolved" concentrations in the water column.

<sup>&</sup>lt;sup>N</sup> The freshwater criterion for copper is a function of the concentration of ions, alkalinity, organic carbon, pH and temperature in the water column. To calculate the criterion, use the Biotic Ligand Model referenced in endnote N at the bottom of Table 30. The acute copper criterion (CMC) is applied as a one-hour average concentration. The chronic criterion (CCC) is applied as a 96-hour (4 days) average concentration. See endnote N also for procedures and information.

				Fresh (µg			water g/L)			
No.	Pollutant	CAS Number	Human Health Criterion	Acute Criterion (CMC)	Chronic Criterion (CCC)	Acute Criterion (CMC)	Chronic Criterion (CCC)			
[Note: The Environmental Quality Commission adopted these revised copper criteria on 11/02/2016.  However, the revised criteria become effective for federal Clean Water Act purposes upon approval by the U.S. Environmental Protection Agency.]										
<u>17</u> 4	Cyanide	57125	у	22 <sup>J</sup>	5.2 <sup>J</sup>	1 <sup>J</sup>	1 <sup>J</sup>			
		<sup>J</sup> This crit	terion is exp	ressed as µg fr	ee cyanide (Cl	V)/L.				
184 5	DDT 4,4'	50293	У	1.1 <sup>A, G</sup>	0.001 <sup>A, G</sup>	0.13 <sup>A, G</sup>	0.001 <sup>A, G</sup>			
A See expanded endnote A at bottom of Table 30 for alternate frequency and duration of this criterion.  G This criterion applies to DDT and its metabolites (i.e. the total concentration of DDT and its metabolites should not exceed this value).										
191 6	Demeton	8065483	n		0.1		0.1			
<u>20</u>	<u>Diazinon</u>	333415	<u>n</u>	0.17	<u>0.17</u>	0.82	0.82			
NOT	E: These diazinon d	criteria are r	ot applicable	e for Clean Wa	ter Act purpose	es until approve	ed by EPA.			
214 7	Dieldrin	60571	у	0.24	0.056	0.71 <sup>A</sup>	0.0019 <sup>A</sup>			
<sup>A</sup> See	e expanded endnot	e A at botto	m of Table 3	30 for alternate	frequency and	duration of this	s criterion.			
<u>22</u> 4 8	Endosulfan	115297	n	0.22 <sup>A , H</sup>	0.056 <sup>A, H</sup>	0.034 <sup>A, H</sup>	0.0087 <sup>A, H</sup>			
A See expanded endnote A at bottom of Table 30 for alternate frequency and duration of this criterion.  H This value is based on the criterion published in Ambient Water Quality Criteria for Endosulfan (EPA 440/5-80-046) and should be applied as the sum of alpha- and beta-endosulfan.										
23 <del>1</del> 9	Endosulfan Alpha	959988	У	0.22 <sup>A</sup>	0.056 A	0.034 A	0.0087 <sup>A</sup>			
<sup>A</sup> See	e expanded endnot	e A at botto	m of Table 3	30 for alternate	frequency and	duration of this	criterion.			

				Fresh			water	
No.	Pollutant	CAS Number	Human Health Criterion	Acute Criterion (CMC)	Chronic Criterion (CCC)	Acute Criterion (CMC)	Chronic Criterion (CCC)	
242 0	Endosulfan Beta	3321365 9	у	0.22 A	0.056 A	0.034 A	0.0087 <sup>A</sup>	
<sup>A</sup> See	e expanded endnot	e A at botto	m of Table 3	30 for alternate	frequency and	duration of this	s criterion.	
25 <del>2</del> 4	Endrin	72208	у	0.086	0.036	0.037 A	0.0023 A	
<sup>A</sup> See	e expanded endnot	e A at botto	m of Table 3	30 for alternate	frequency and	duration of this	s criterion.	
<u>26</u> 2	Guthion	86500	n		0.01		0.01	
<u>27</u> 2 3	Heptachlor	76448	у	0.52 <sup>A</sup>	0.0038 A	0.053 A	0.0036 A	
<sup>A</sup> See	e expanded endnot	e A at botto	m of Table 3	30 for alternate	frequency and	duration of this	s criterion.	
<u>28</u> 2 4	Heptachlor Epoxide	1024573	у	0.52 <sup>A</sup>	0.0038 A	0.053 <sup>A</sup>	0.0036 A	
<sup>A</sup> See	e expanded endnot	e A at botto	m of Table 3	30 for alternate	frequency and	duration of this	s criterion.	
<u>29</u> 2 5	Iron (total)	7439896	n		1000			
30 <del>2</del> 6	Lead	7439921	n	See C, F	See C, F	210 <sup>c</sup>	8.1 <sup>c</sup>	
<sup>c</sup> Criterion is expressed in terms of "dissolved" concentrations in the water column. <sup>F</sup> The freshwater criterion for this metal is expressed as a function of hardness (mg/L) in the water column.  To calculate the criterion, use formula under expanded endnote F at bottom of Table 30.								
31 <sub>2</sub> 7	Malathion	121755	n		0.1		0.1	
32 <del>2</del> 8	Mercury (total)	7439976	n	2.4	0.012	2.1	0.025	

					Freshwater (μg/L)		twater ug/L)
No.	Pollutant	CAS Number	Human Health Criterion	Acute Criterion (CMC)	Chronic Criterion (CCC)	Acute Criterion (CMC)	Chronic Criterion (CCC)
33 <del>2</del> 9	Methoxychlor	72435	у	1	0.03	1	0.03
343 0	Mirex	2385855	n	-	0.001	-	0.001
35 <del>3</del> 4	Nickel	7440020	у	See C, F	See C, F	74 <sup>c</sup>	8.2 <sup>c</sup>
F The	<sup>c</sup> Criterion is expressed in terms of "dissolved" concentrations in the water column. <sup>F</sup> The freshwater criterion for this metal is expressed as a function of hardness (mg/L) in the water column.  To calculate the criterion, use formula under expanded endnote F at bottom of Table 30.						
36 <del>3</del> 2	Parathion	56382	n	0.065	0.013	-	
373 3	Pentachlorophe nol	87865	у	See I	See I	13	7.9
	shwater aquatic life llows: CMC=(exp(1	•	-	•		ction of pH, an	d are calculated
38 <del>3</del> 4	Phosphorus Elemental	7723140	n				0.1
393 5	Polychlorinated Biphenyls (PCBs)	NA	У	2 <sup>K</sup>	0.014 <sup>K</sup>	10 <sup>K</sup>	0.03 <sup>K</sup>
K Thi	s criterion applies to	o total PCBs	s (e.g. deterr	nined as Arock	ors or congene	rs)	
<u>40</u> 3 6	Selenium	7782492	у	See C , L	4.6 <sup>c</sup>	290 <sup>c</sup>	71 <sup>C</sup>

<sup>&</sup>lt;sup>c</sup> Criterion is expressed in terms of "dissolved" concentrations in the water column.

L The CMC=(1/[(f1/CMC1)+(f2/CMC2)]μg/L) \* CF where f1 and f2 are the fractions of total selenium that are treated as selenite and selenate, respectively, and CMC1 and CMC2 are 185.9 μg/L and 12.82 μg/L, respectively. See expanded endnote F for the Conversion Factor (CF) for selenium.

				Fresh (µg		Saltwater (µg/L)		
No.	Pollutant	CAS Number	Human Health Criterion	Acute Criterion (CMC)	Chronic Criterion (CCC)	Acute Criterion (CMC)	Chronic Criterion (CCC)	
413 7	Silver	7440224	n	See C , F	0.10 <sup>c</sup>	1.9 <sup>c</sup>		

<sup>&</sup>lt;sup>c</sup> Criterion is expressed in terms of "dissolved" concentrations in the water column.

F The freshwater acute criterion for this metal is expressed as a function of hardness (mg/L) in the water column. To calculate the criterion, use formula under expanded endnote F at bottom of Table 30.

0 0 1 0 11	ordanim to dalcalate the officinent, add formala and of expanded officine to account of facile del									
<u>42</u> 3 8	Sulfide Hydrogen Sulfide	7783064	n		2	l	2			
433 9	Toxaphene	8001352	у	0.73	0.0002	0.21	0.0002			
444 0	Tributyltin (TBT)	688733	n	0.46	0.063 0.072	<del>0.37</del> <u>0.42</u>	<del>0.01</del> <u>0.0074</u>			

NOTE: The freshwater chronic and saltwater acute and chronic criteria for tributyltin are not applicable for Clean Water Act purposes until the revised Table 30 values are approved by EPA. The freshwater acute criterion is not changing and is applicable. The following tributyltin criteria remain applicable until EPA approves the revised criteria shown in Table 30: the freshwater chronic (CCC) is 0.063 µg/L, the saltwater acute (CMC) is 0.37 µg/L, and the saltwater chronic (CCC) is 0.01 µg/L.

<u>45</u> 4	Zinc	7440666	у	See C, F	See C , F	90 c	81 <sup>c</sup>
4							

<sup>&</sup>lt;sup>c</sup> Criterion is expressed in terms of "dissolved" concentrations in the water column.

Expanded Endnotes A, E, F, N, O

F The freshwater criterion for this metal is expressed as a function of hardness (mg/L) in the water column. To calculate the criterion, use formula under expanded endnote F at bottom of Table 30.

### **Endnote A: Alternate Frequency and Duration for Certain Pesticides**

This criterion is based on EPA recommendations issued in 1980 that were derived using guidelines that differed from EPA's 1985 Guidelines which update minimum data requirements and derivation procedures. The CMC may not be exceeded at any time and the CCC may not be exceeded based on a 24-hour average. The CMC may be applied using a one hour averaging period not to be exceeded more than once every three years, if the CMC values given in Table 30 are divided by 2 to obtain a value that is more comparable to a CMC derived using the 1985 Guidelines.

### **Endnote E: Equation for Hardness-Dependent Freshwater Cadmium Acute Criteria**

The freshwater criterion for this metal is expressed as total recoverable with two significant figures, and is a function of hardness (mg/L) in the water column. Criteria values based on hardness are calculated using the following formula (CMC refers to the acute criterion):

**CMC** = 
$$(exp(m_A*[ln(hardness)] + b_A))$$

Chemical	<b>m</b> ₄	<b>b</b> ∧	mc	<del>b</del> c
Cadmium	<del>1.128</del>	<del>-3.828</del>	N/A	<del>N/A</del>

### **Endnote F: Equations for Hardness-Dependent Freshwater Metals Criteria and Conversion Factor Table**

The freshwater criterion for this metal is expressed as dissolved with two significant figures, and is a function of hardness (mg/L) in the water column. Criteria values based on hardness are calculated using the following formulas (CMC refers to the acute criterion; CCC refers to the chronic criterion):

**CMC** = 
$$(exp(m_A*[ln(hardness)] + b_A))*CF f$$

**CCC** = 
$$(exp(m_C*[ln(hardness)] + b_C))*CF$$

"CF" is the conversion factor used for converting a metal criterion expressed as the total recoverable fraction in the water column to a criterion expressed as the dissolved fraction in the water column.

Values for Calcu	lating Hardno	ess-Depende	nt Metals Cri	teria								
Chemical m <sub>A</sub> b <sub>A</sub> m <sub>C</sub> b <sub>C</sub>												
Cadmium	0.9789 N/A	<u>-3.866</u> N/A	0.7409	-4.719								
Chromium III	0.8190	3.7256	0.8190	0.6848								
Lead	1.273	-1.460	1.273	-4.705								
Nickel	0.8460	2.255	0.8460	0.0584								
Silver	1.72	-6.59										
Zinc	0.8473	0.884	0.8473	0.884								

The conversion factors (CF) below must be used in the equations above for the hardness-dependent metals in order to convert total recoverable metals criteria to dissolved metals criteria. For metals that are not hardness-dependent (i.e. arsenic, chromium VI, selenium, and silver (chronic)), or are saltwater criteria, the criterion value associated with the metal in Table 30 already reflects a dissolved criterion based on its conversion factor below.

	Conversion Factor (	CF) Table for Dissolv	ed Metals			
Chemical	Fresh	water	Sa	ltwater		
Chemical	Acute	Chronic	Acute	Chronic		
Arsenic	1.000	1.000	1.000	1.000		
Cadmium	1.136672-[(ln hardness)(0.041838)] N/A	1.101672-[(ln hardness)(0.041838)]	0.994	0.994		
Chromium III	0.316	0.860				
Chromium VI	0.982	0.962	0.993	0.993		
Copper	N/A	N/A	0.83	0.83		
Lead	1.46203-[(In hardness)(0.145712)]	1.46203-[(In hardness)(0.145712)]	0.951	0.951		
Nickel	0.998	0.997	0.990	0.990		
Selenium	0.996	0.922	0.998	0.998		
Silver	0.85	0.85	0.85			
Zinc	0.978	0.986	0.946	0.946		

NOTE: Cadmium values  $m_A$ ,  $b_A$ , and the freshwater acute CF in Endnote F are not applicable until the freshwater acute cadmium criterion in Table 30 is approved by EPA. Until then the federally promulgated freshwater acute cadmium criterion at 40 CFR 131.46 remains applicable for CWA purposes.

### **Endnote N: Deriving freshwater copper criteria**

The freshwater copper criteria at any time are the Biotic Ligand Model (BLM) derived Instantaneous Water Quality Criteria (IWQC) output based on a concurrently measured set of model input parameter values. The Biotic Ligand Model uses multiple ambient water quality parameters to derive 1-hour acute exposure (CMC) and 96-hour chronic exposure (CCC) water quality criteria (IWQC) for copper based on the site specific water chemistry that determines the toxicity of copper to aquatic life. If measured data for one or more of the model input parameters used to derive the acute and chronic IWQC is not available, the procedures in section (1) or (2) of this endnote will be used as specified to substitute an estimate or a default value for the missing input parameter. BLM results (IWQC) based on sufficient measured input parameter data are more accurate and supersede results based on estimates or default values. The acceptable BLM software to calculate the IWQC include version 2.2.3, referenced in "Aquatic Life Ambient Freshwater Quality Criteria – Copper": EPA-822-R-07-001, February 2007, and version 2.2.4. The criteria are expressed as dissolved copper in micrograms per liter (to the nearest one-tenth).

(1) Input Parameter Substitution and Estimation Procedures to Derive BLM Criteria (IWQC)

If the measured value for any input parameter needed to derive an IWQC using the BLM is not available, DEQ will substitute an estimated input parameter value according to the procedures described in this section [Endnote N (1)]. If the data required to determine the estimated parameter value is not available, DEQ will use default values derived according to the procedures in Endnote N (2).

- (a) Total recoverable concentration measurements will be substituted for dissolved concentration measurements that are not available. For alkalinity, calcium, chloride, magnesium, potassium, sodium and sulfate, total recoverable concentration measurements will be used as a direct substitute for dissolved concentration measurements. Total organic carbon (TOC) measurements will be multiplied by 0.83 to convert the TOC value to an equivalent dissolved organic carbon (DOC) value; except where sufficient TOC and DOC data are available for a site, DEQ will calculate and apply a site-specific translator in place of 0.83 to convert TOC values to DOC for use in the BLM.
- (b) Alkalinity, calcium, chloride, magnesium, potassium, sodium and sulfate: If data for any of these BLM input parameters are missing from a particular dataset, DEQ will estimate its value based on the relationship of the ion or alkalinity to specific conductance measurements for that data set using the regression analysis equations in Table 1. Specific conductance measurements must be concurrent with the other BLM input parameters dataset.

	Table N-1
Parameter	Regression Equation
Alkalinity	Alk. = $\exp^{(0.88 \cdot [\ln(SpC)] - 0.41)}$
Calcium	Ca = $\exp^{(0.96 \cdot [\ln(SpC)] - 2.29)}$
Chloride	$CI = exp^{(1.15\cdot[ln(SpC)] - 3.82)}$
Magnesium	$Mg = exp^{(0.91 \cdot [ln(SpC)] - 3.09)}$
Potassium	$K = exp^{(0.84 \cdot [ln(SpC)] - 3.74)}$
Sodium	Na = $\exp^{(0.86 \cdot [\ln(SpC)] - 2.22)}$
Sulfate	$SO_4 = exp^{(1.45 \cdot [ln(SpC)] - 5.59}$

Where, "SpC" is a measurement of specific conductance in µmhos/cm, "In" is the natural logarithm, and "exp" is a mathematical constant that is the base of the natural logarithm.

### (c) pH

If concurrent pH data is missing from the sample dataset, DEQ will use a representative pH value determined by interpolating from data available for the site or proximate monitoring locations where conditions (such as type of water body, stream flow and geology) are similar to the site. DEQ will use the available data and methods to produce the best practicable estimate of pH for the site and time for which the IWQC is being derived.

### (d) Temperature

If concurrent temperature data is missing from the sample dataset, DEQ will use a monthly mean temperature based on data available for the site or proximate monitoring locations where conditions (such as type of water body and stream flow) are similar to the site.

### (e) Humic Acid

If sufficient high quality data on the percentage of humic acid as a proportion of DOC is available for a site, DEQ will use that value in the BLM in place of the default value of 10% used in the model.

### (2) Default Action Values

If the measured value for DOC, alkalinity, calcium, chloride, magnesium, potassium, sodium or sulfate is not available to derive an IWQC using the BLM, and the parameter value cannot be estimated as specified in section (1) above, DEQ will use a conservative input value for the missing parameter as described in this section [Endnote N (2)] to derive a default action value using the Biotic Ligand Model. The default action value will be used for Clean Water Act purposes until measured or estimated input parameter data are available to derive accurate copper criteria (IWQC) based on site specific water chemistry.

(a) The default input parameter values for DOC, alkalinity calcium, chloride, magnesium, potassium, sodium and sulfate will be the percentile value from the

distribution of the high quality data available for surface waters in the region as shown in Table N-2.

	Table N-2 data distributio fault value by re	n to be used as egion													
Region	Region DOC percentile Alkalinity and lons percentile														
Willamette	20 <sup>th</sup>	20 <sup>th</sup>													
Coastal	20 <sup>th</sup>	20 <sup>th</sup>													
Cascades	20 <sup>th</sup>	20 <sup>th</sup>													
Eastern	15 <sup>th</sup>	15 <sup>th</sup>													
Columbia River	20 <sup>th</sup>	20 <sup>th</sup>													

- (b) The regional default values for each parameter and region will be updated periodically as additional high quality data becomes available and is added to DEQ's database.
- (c) The regional default values for each parameter are available on DEQ's website.
- (d) The regions listed in Table N-2 are comprised of the following EPA Level III ecoregions or waterbody:
  - (i) Willamette: the Willamette Valley
  - (ii) Coastal: Coast Range and Klamath Mountains
  - (iii) Cascades: Cascades
  - (iv) Eastern: Eastern Cascades Slopes and Foothills, Columbia Plateau, Blue

Mountains, Northern Basin and Range and Snake River Plain

(v) Columbia River: Columbia River mainstem in Oregon

### (3) General Policies

- (a) The copper BLM derives instantaneous criteria results (IWQC) that vary at a site over time reflecting the effect of local water chemistry on copper toxicity to aquatic organisms. DEQ will apply the BLM criteria for Clean Water Act purposes to protect the water body during the most bioavailable or toxic conditions.
- (b) For assessing waters of the state, DEQ will use approaches that give preference to the use of BLM criteria derived with site-specific measured input parameter data.

### **Endnote O: Deriving freshwater aluminum criteria**

Freshwater aluminum criteria values are calculated from a concurrently measured set of Aluminum Criteria Calculator (ACC) input parameter values. The input parameter values are based on the site and time specific water chemistry that determines the toxicity of aluminum to aquatic life. If measured data for one or more of the ACC input parameters is not available, the procedures in section (1), (2), or (3) of this endnote will be used, in the order listed, as specified to substitute an estimated or a default value for the missing input parameter or to apply default criteria derived using ecoregional data.

Criteria values based on sufficient concurrent measured input parameter data are more accurate, preferred, and supersede results based on estimates, default values or applied default ecoregional criteria values. The criteria are expressed as total recoverable in micrograms per liter (to two significant figures). Oregon will use analytical methods that measure the bioavailable fraction of aluminum unless total recoverable aluminum measurements are required by Federal regulations.

### (1) Input Parameter Estimation Procedures to Derive Aluminum Criteria Values

If the measured value for one or more input parameters needed to derive an aluminum criteria value using the ACC is not available, DEQ will substitute a calculated or estimated input value according to the procedures described in this section [Endnote O (1)].

### (a) DOC

DEQ will use total organic carbon (TOC) measurements to estimate DOC measurements that are not available. Total organic carbon (TOC) measurements will be multiplied by 0.83 to convert the TOC value to an equivalent dissolved organic carbon (DOC) value; except where sufficient TOC and DOC data are available for a site, DEQ will calculate and apply a site-specific translator in place of 0.83 to convert TOC values to DOC for use in the Aluminum Criteria Calculator. If neither DOC nor TOC measurements are available, substitute a default DOC value as described in Endnote O (2).

### (b) Total Hardness

If total hardness is not available, DEQ will estimate total hardness by substituting dissolved hardness as an input parameter for the Aluminum Criteria Calculator. If neither total nor dissolved hardness data are available, DEQ will use the equation in Table O-1 to estimate total hardness using specific conductance. Specific conductance measurements must be concurrent with the other input parameters for the Aluminum Criteria Calculator. If total hardness cannot be estimated from concurrent data, DEQ will apply the applicable ecoregional default aluminum criterion described in Endnote O (3).

	Table O-1 estimate total hardness from ecific conductance
<u>Parameter</u>	Regression Equation
Total Hardness	$\underline{\text{Total Hardness} = \exp^{(1.050 \cdot [\ln(\text{SpC})] - 1.211)}}$

Where, "SpC" is a measurement of specific conductance in µmhos/cm, "In" is the natural logarithm, and "exp" is a mathematical constant that is the base of the natural logarithm.

### (2) Applying a Default Value for DOC to Derive Aluminum Criteria Values

If concurrently measured DOC is not available to derive a criteria value using the ACC and DOC cannot be estimated as specified in Endnote O (1)(a) above, DEQ will use a conservative

default DOC input value as described in this section [Endnote O (2)] to derive the criteria value. The default DOC input value will be used for Clean Water Act purposes until measured or estimated DOC input data are available to derive aluminum criteria values based on site-specific water chemistry.

(a) The default input parameter values for DOC will be the percentile value from the distribution of the high-quality data available for surface waters in the region as shown in Table O-2.

Percentile of data	e O-2 distribution to be value by region
Region	<u>DOC</u> <u>percentile</u>
Willamette	<u>15<sup>th</sup></u>
Coastal	<u>30<sup>th</sup></u>
Cascades	<u>20<sup>th</sup></u>
<u>Eastern</u>	<u>15<sup>th</sup></u>
Columbia River	<u>10<sup>th</sup></u>

- b) The regional default DOC values will be updated periodically as additional high-quality data become available and are added to DEQ's database.
- (c) The resulting regional default input values for DOC are shown on DEQ's website.
- (d) The regions listed in Table O-2 are the same as those defined in Endnote N(2)(d).
- (3) Applying Aluminum Default Ecoregional Criteria

If data for pH is missing or hardness is missing and cannot be estimated as described in Endnote O (1)(b), DEQ will apply an ecoregional default aluminum criteria value.

- (a) The default ecoregional acute (CMC) and chronic (CCC) criteria values will be the 10<sup>th</sup> percentile value from the distribution of all ACC outputs calculated from concurrently measured high quality input data available for Oregon surface waters by EPA Level III ecoregion with the Columbia River mainstem treated separately.
- (b) The ecoregional default aluminum criteria values will be updated periodically as additional high quality data become available and are added to DEQ's database.
- (c) The resulting ecoregional default aluminum criteria values are shown on DEQ's website.

### (4) General Policies

- (a) The ACC produces criteria values that vary at a site over time reflecting the effect of local water chemistry on aluminum toxicity to aquatic organisms. To apply the aluminum criteria for Clean Water Act purposes, criteria values based on ambient water chemistry conditions must protect the water body over the full range of water chemistry conditions, including during conditions when aluminum is most toxic.
- (b) When applying the aluminum criteria, DEQ will use approaches that give preference to the use of ACC criteria values based on concurrently measured or estimated input parameter data, as described in Endnote O (1), (2), and (3) in the order listed, and concurrently measured aluminum data.

### **Table 30(a): Ammonia Acute Criteria Values (One-hour Average)**—Salmonid Species Present Temperature and pH-Dependent and expressed as Total Ammonia Nitrogen (mg/L TAN)

### Criteria cannot be exceeded more than once every three years

$$Acute \ Criterion = MIN \left( \left( \frac{0.275}{1 + 10^{7.204 - pH}} + \frac{39.0}{1 + 10^{pH - 7.204}} \right), \left( 0.7249 \times \left( \frac{0.0114}{1 + 10^{7.204 - pH}} + \frac{1.6181}{1 + 10^{pH - 7.204}} \right) \times \left( 23.12 \times 10^{0.036 \times (20 - T)} \right) \right) \right)$$

							Т	empera	iture (°C	:)							/ /
рН	0-14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
6.5	33	33	32	29	27	25	23	21	19	18	16	15	14	13	12	11	9.9
6.6	31	31	30	28	26	24	22	20	18	17	16	14	13	12	11	10	9.5
6.7	30	30	29	27	24	22	21	19	18	16	15	14	13	12	11	9.8	9.0
6.8	28	28	27	25	23	21	20	18	17	15	14	13	12	11	10	9.2	8.5
6.9	26	26	25	23	21	20	18	17	15	14	13	12	11	10	9.4	8.6	7.9
7.0	24	24	23	21	20	18	17	15	14	13	12	11	10	9.4	8.6	8.0	7.3
7.1	22	22	21	20	18	17	15	14	13	12	11	10	9.3	8.5	7.9	7.2	6.7
7.2	20	20	19	18	16	15	14	13	12	11	9.8	9.1	8.3	7.7	7.1	6.5	6.0
7.3	18	18	17	16	14	13	12	11	10	9.5	8.7	8.0	7.4	6.8	6.3	5.8	5.3
7.4	15	15	15	14	13	12	11	9.8	9.0	8.3	7.7	7.0	6.5	6.0	5.5	5.1	4.7
7.5	13	13	13	12	11	10	9.2	8.5	7.8	7.2	6.6	6.1	5.6	5.2	4.8	4.4	4.0
7.6	11	11	11	10	9.3	8.6	7.9	7.3	6.7	6.2	5.7	5.2	4.8	4.4	4.1	3.8	3.5
7.7	9.6	9.6	9.3	8.6	7.9	7.3	6.7	6.2	5.7	5.2	4.8	4.4	4.1	3.8	3.5	3.2	3.0
7.8	8.1	8.1	7.9	7.2	6.7	6.1	5.6	5.2	4.8	4.4	4.0	3.7	3.4	3.2	2.9	2.7	2.5
7.9	6.8	6.8	6.6	6.0	5.6	5.1	4.7	4.3	4.0	3.7	3.4	3.1	2.9	2.6	2.4	2.2	2.1
8.0	5.6	5.6	5.4	5.0	4.6	4.2	3.9	3.6	3.3	3.0	2.8	2.6	2.4	2.2	2.0	1.9	1.7
8.1	4.6	4.6	4.5	4.1	3.8	3.5	3.2	3.0	2.7	2.5	2.3	2.1	2.0	1.8	1.7	1.5	1.4
8.2	3.8	3.8	3.7	3.5	3.1	2.9	2.7	2.4	2.3	2.1	1.9	1.8	1.6	1.5	1.4	1.3	1.2
8.3	3.1	3.1	3.1	2.8	2.6	2.4	2.2	2.0	1.9	1.7	1.6	1.4	1.3	1.2	1.1	1.0	0.96
8.4	2.6	2.6	2.5	2.3	2.1	2.0	1.8	1.7	1.5	1.4	1.3	1.2	1.1	1.0	0.93	0.86	0.79
8.5	2.1	2.1	2.1	1.9	1.8	1.6	1.5	1.4	1.3	1.2	1.1	0.98	0.90	0.83	0.77	0.71	0.65
8.6	1.8	1.8	1.7	1.6	1.5	1.3	1.2	1.1	1.0	0.96	0.88	0.81	0.75	0.69	0.63	0.59	0.54
8.7	1.5	1.5	1.4	1.3	1.2	1.1	1.0	0.94	0.87	0.80	0.74	0.68	0.62	0.57	0.53	0.49	0.45
8.8	1.2	1.2	1.2	1.1	1.0	0.93	0.86	0.79	0.73	0.67	0.62	0.57	0.52	0.48	0.44	0.41	0.37
8.9	1.0	1.0	1.0	0.93	0.85	0.79	0.72	0.67	0.61	0.56	0.52	0.48	0.44	0.40	0.37	0.34	0.32
9.0	0.88	0.88	0.86	0.79	0.73	0.67	0.62	0.57	0.52	0.48	0.44	0.41	0.37	0.34	0.32	0.29	0.27

### Table 30(b): Ammonia Acute Criteria Values (One-hour Average\*)—Salmonid Species Absent

Temperature and pH-Dependent and expressed as Total Ammonia Nitrogen (mg/L TAN)

### Criteria cannot be exceeded more than once every three years

Acute Criterion =  $0.7249 \times \frac{0.0114}{1 + 10^{7.204 - pH}} + \frac{1.6181}{1 + 10^{pH - 7.204}} \times MIN(51.93, 23.12 \times 10^{0.036 \times (20 - T)})$ Temperature (°C)

												<b>O</b> )									
рН	0-10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
6.5	51	48	44	41	37	34	32	29	27	25	23	21	19	18	16	15	14	13	12	11	9.9
6.6	49	46	42	39	36	33	30	28	26	24	22	20	18	17	16	14	13	12	11	10	9.5
6.7	46	44	40	37	34	31	29	27	24	22	21	19	18	16	15	14	13	12	11	9.8	9.0
6.8	44	41	38	35	32	30	27	25	23	21	20	18	17	15	14	13	12	11	10	9.2	8.5
6.9	41	38	35	32	30	28	25	23	21	20	18	17	15	14	13	12	11	10	9.4	8.6	7.9
7.0	38	35	33	30	28	25	23	21	20	18	17	15	14	13	12	11	10	9.4	8.6	7.9	7.3
7.1	34	32	30	27	25	23	21	20	18	17	15	14	13	12	11	10	9.3	8.5	7.9	7.2	6.7
7.2	31	29	27	25	23	21	19	18	16	15	14	13	12	11	9.8	9.1	8.3	7.7	7.1	6.5	6.0
7.3	27	26	24	22	20	18	17	16	14	13	12	11	10	9.5	8.7	8.0	7.4	6.8	6.3	5.8	5.3
7.4	24	22	21	19	18	16	15	14	13	12	11	9.8	9.0	8.3	7.7	7.0	6.5	6.0	5.5	5.1	4.7
7.5	21	19	18	17	15	14	13	12	11	10	9.2	8.5	7.8	7.2	6.6	6.1	5.6	5.2	4.8	4.4	4.0
7.6	18	17	15	14	13	12	11	10	9.3	8.6	7.9	7.3	6.7	6.2	5.7	5.2	4.8	4.4	4.1	3.8	3.5
7.7	15	14	13	12	11	10	9.3	8.6	7.9	7.3	6.7	6.2	5.7	5.2	4.8	4.4	4.1	3.8	3.5	3.2	2.9
7.8	13	12	11	10	9.3	8.5	7.9	7.2	6.7	6.1	5.6	5.2	4.8	4.4	4.0	3.7	3.4	3.2	2.9	2.7	2.5
7.9	11	9.9	9.1	8.4	7.7	7.1	6.6	3.0	5.6	5.1	4.7	4.3	4.0	3.7	3.4	3.1	2.9	2.6	2.4	2.2	2.1
8.0	8.8	8.2	7.6	7.0	6.4	5.9	5.4	5.0	4.6	4.2	3.9	3.6	3.3	3.0	2.8	2.6	2.4	2.2	2.0	1.9	1.7
8.1	7.2	6.8	6.3	5.8	5.3	4.9	4.5	4.1	3.8	3.5	3.2	3.0	2.7	2.5	2.3	2.1	2.0	1.8	1.7	1.5	1.4
8.2	6.0	5.6	5.2	4.8	4.4	4.0	3.7	3.4	3.1	2.9	2.7	2.4	2.3	2.1	1.9	1.8	1.6	1.5	1.4	1.3	1.2
8.3	4.9	4.6	4.3	3.9	3.6	3.3	3.1	2.8	2.6	2.4	2.2	2.0	1.9	1.7	1.6	1.4	1.3	1.2	1.1	1.0	0.96
8.4	4.1	3.8	3.5	3.2	3.0	2.7	2.5	2.3	2.1	2.0	1.8	1.7	1.5	1.4	1.3	1.2	1.1	1.0	0.93	0.86	0.79
8.5	3.3	3.1	2.9	2.7	2.4	2.3	2.1	1.9	1.8	1.6	1.5	1.4	1.3	1.2	1.1	0.98	0.90	0.83	0.77	0.71	0.65
8.6	2.8	2.6	2.4	2.2	2.0	1.9	1.7	1.6	1.5	1.3	1.2	1.1	1.0	0.96	0.88	0.81	0.75	0.69	0.63	0.58	0.54
8.7	2.3	2.2	2.0	1.8	1.7	1.6	1.4	1.3	1.2	1.1	1.0	0.94	0.87	0.80	0.74	0.68	0.62	0.57	0.53	0.49	0.45
8.8	1.9	1.8	1.7	1.5	1.4	1.3	1.2	1.1	1.0	0.93	0.86	0.79	0.73	0.67	0.62	0.57	0.52	0.48	0.44	0.41	0.37
8.9	1.6	1.5	1.4	1.3	1.2	1.1	1.0	0.93	0.85	0.79	0.72	0.67	0.61	0.56	0.52	0.48	0.44	0.40	0.37	0.34	0.32
9.0	1.4	1.3	1.2	1.1	1.0	0.93	0.86	0.79	0.73	0.67	0.62	0.57	0.52	0.48	0.44	0.41	0.37	0.34	0.32	0.29	0.27

### Table 30(c): Ammonia Chronic Criteria Values (30-day Rolling Average\*)

Temperature and pH-Dependent and expressed as Total Ammonia Nitrogen (mg/L TAN)

\* The highest four-day average within the 30-day averaging period must not be more than 2.5 times the chronic value

### Criteria cannot be exceeded more than once every three years

Chronic Criterion = 
$$0.8876 \times \left(\frac{0.0278}{1 + 10^{7.688 - pH}} + \frac{1.1994}{1 + 10^{pH - 7.688}}\right) \times \left(2.126 \times 10^{0.028 \times \left(20 - MAX(T,7)\right)}\right)$$

									\ <u>1</u>	+ 10'		eratur		7.000	,									
рН	0-7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
6.5	4.9	4.6	4.3	4.1	3.8	3.6	3.3	3.1	2.9	2.8	2.6	2.4	2.3	2.1	2.0	1.9	1.8	1.6	1.5	1.5	1.4	1.3	1.2	1.1
6.6	4.8	4.5	4.3	4.0	3.8	3.5	3.3	3.1	2.9	2.7	2.5	2.4	2.2	2.1	2.0	1.8	1.7	1.6	1.5	1.4	1.3	1.3	1.2	1.1
6.7	4.8	4.5	4.2	3.9	3.7	3.5	3.2	3.0	2.8	2.7	2.5	2.3	2.2	2.1	1.9	1.8	1.7	1.6	1.5	1.4	1.3	1.2	1.2	1.1
6.8	4.6	4.4	4.1	3.8	3.6	3.4	3.2	3.0	2.8	2.6	2.4	2.3	2.1	2.0	1.9	1.8	1.7	1.6	1.5	1.4	1.3	1.2	1.1	1.1
6.9	4.5	4.2	4.0	3.7	3.5	3.3	3.1	2.9	2.7	2.5	2.4	2.2	2.1	2.0	1.8	1.7	1.6	1.5	1.4	1.3	1.2	1.2	1.1	1.0
7.0	4.4	4.1	3.8	3.6	3.4	3.2	3.0	2.8	2.6	2.4	2.3	2.2	2.0	1.9	1.8	1.7	1.6	1.5	1.4	1.3	1.2	1.1	1.1	0.99
7.1	4.2	3.9	3.7	3.5	3.2	3.0	2.8	2.7	2.5	2.3	2.2	2.1	1.9	1.8	1.7	1.6	1.5	1.4	1.3	1.2	1.2	1.1	1.0	0.95
7.2	4.0	3.7	3.5	3.3	3.1	2.9	2.7	2.5	2.4	2.2	2.1	2.0	1.8	1.7	1.6	1.5	1.4	1.3	1.3	1.2	1.1	1.0	0.96	0.90
7.3	3.8	3.5	3.3	3.1	2.9	2.7	2.6	2.4	2.2	2.1	2.0	1.8	1.7	1.6	1.5	1.4	1.3	1.3	1.2	1.1	1.0	0.97	0.91	0.85
7.4	3.5	3.3	3.1	2.9	2.7	2.5	2.4	2.2	2.1	2.0	1.8	1.7	1.6	1.5	1.4	1.3	1.3	1.2	1.1	1.0	0.96	0.90	0.85	0.79
7.5	3.2	3.0	2.8	2.7	2.5	2.3	2.2	2.1	1.9	1.8	1.7	1.6	1.5	1.4	1.3	1.2	1.2	1.1	1.0	0.95	0.89	0.83	0.78	0.73
7.6	2.9	2.8	2.6	2.4	2.3	2.1	2.0	1.9	1.8	1.6	1.5	1.4	1.4	1.3	1.2	1.1	1.1	0.98	0.92	0.86	0.81	0.76	0.71	0.67
7.7	2.6	2.4	2.3	2.2	2.0	1.9	1.8	1.7	1.6	1.5	1.4	1.3	1.2	1.1	1.1	1.0	0.94	0.88	0.83	0.78	0.73	0.68	0.64	0.60
7.8	2.3	2.2	2.1	1.9	1.8	1.7	1.6	1.5	1.4	1.3	1.2	1.2	1.1	1.0	0.95	0.89	0.84	0.79	0.74	0.69	0.65	0.61	0.57	0.53
7.9	2.1	1.9	1.8	1.7	1.6	1.5	1.4	1.3	1.2	1.2	1.1	1.0	0.95	0.89	0.84	0.79	0.74	0.69	0.65	0.61	0.57	0.53	0.50	0.47
8.0	1.8	1.7	1.6	1.5	1.4	1.3	1.2	1.1	1.1	1.0	0.94	0.88	0.83	0.78	0.73	0.68	0.64	0.60	0.56	0.53	0.50	0.44	0.44	0.41
8.1	1.5	1.5	1.4	1.3	1.2	1.1	1.1	0.99	0.92	0.87	0.81	0.76	0.71	0.67	0.63	0.59	0.55	0.52	0.49	0.46	0.43	0.40	0.38	0.35
8.2	1.3	1.2	1.2	1.1	1.0	0.96	0.90	0.84	0.79	0.74	0.70	0.65	0.61	0.57	0.54	0.50	0.47	0.44	0.42	0.39	0.37	0.34	0.32	0.30
8.3	1.1	1.1	0.99	0.93	0.87	0.82	0.76	0.72	0.67	0.63	0.59	0.55	0.52	0.49	0.46	0.43	0.40	0.38	0.35	0.33	0.31	0.29	0.27	0.26
8.4	0.95	0.89	0.84	0.79	0.74	0.69	0.65	0.61	0.57	0.53	0.50	0.47	0.44	0.41	0.39	0.36	0.34	0.32	0.30	0.28	0.26	0.25	0.23	0.22
8.5	0.80	0.75	0.71	0.67	0.62	0.58	0.55	0.51	0.48	0.45	0.42	0.40	0.37	0.35	0.33	0.31	0.29	0.27	0.25	0.24	0.22	0.21	0.20	0.18
8.6	0.68	0.64	0.60	0.56	0.53	0.49	0.46	0.43	0.41	0.38	0.36	0.33	0.31	0.29	0.28	0.26	0.24	0.23	0.21	0.20	0.19	0.18	0.16	0.15
8.7	0.57	0.54	0.51	0.47	0.44	0.42	0.39	0.37	0.34	0.32	0.30	0.28	0.27	0.25	0.23	0.22	0.21	0.19	0.18	0.17	0.16	0.15	0.14	0.13
8.8	0.49	0.46	0.43	0.40	0.38	0.35	0.33	0.31	0.29	0.27	0.26	0.24	0.23	0.21	0.20	0.19	0.17	0.16	0.15	0.14	0.13	0.13	0.12	0.11
8.9	0.42	0.39	0.37	0.34	0.32	0.30	0.28	0.27	0.25	0.23	0.22	0.21	0.19	0.18	0.17	0.16	0.15	0.14	0.13	0.12	0.12	0.11	0.10	0.09
9.0	0.36	0.34	0.32	0.30	0.28	0.26	0.24	0.23	0.21	0.20	0.19	0.18	0.17	0.16	0.15	0.14	0.13	0.12	0.11	0.11	0.10	0.09	0.09	0.08



### Effective April 18, 2014

### Water Quality Guidance Values Summary A

The concentration for each compound listed in Table 31 is a guidance value that DEQ may use in application of Oregon's Toxic Substances Narrative (340-041-0033(2)) to waters of the state in order to protect aquatic life. All values are expressed as micrograms per liter (µg/L) except where noted. Compounds are listed in alphabetical order with the corresponding EPA number (from National Recommended Water Quality Criteria: 2002, EPA-822-R-02-047), corresponding Chemical Abstract Service (CAS) number, aquatic life freshwater acute and chronic guidance values, and aquatic life saltwater acute and chronic guidance values.

### OAR 340-041-8033 Table 31 Aquatic Life Water Quality Guidance Values for Toxic Pollutants

EPA	Dellestant	CAS	Fresh	water	Salt	water
No.	<del>Pollutant</del>	CAS Number	Acute	Chronic	Acute	Chronic
<del>56</del>	Acenaphthene	83329	<del>1,700</del>	<del>520</del>	970	<del>710</del>
<del>17</del>	Acrolein	107028	<del>68</del>	<del>21</del>	<del>55</del>	
<del>18</del>	Acrylonitrile	107131	<del>7,550</del>	<del>2,600</del>		
4	Antimony	7440360	9,000	<del>1,600</del>		
<del>19</del>	Benzene	<del>71432</del>	<del>5,300</del>		<del>5,100</del>	<del>700</del>
<del>59</del>	Benzidine	92875	<del>2,500</del>			
3	Beryllium	7440417	<del>130</del>	<del>5.3</del>		
19 B	BHC (Hexachlerocyclohexa ne-Technical)	319868	<del>100</del>		0.34	
<del>21</del>	Carbon Tetrachloride	<del>56235</del>	<del>35,200</del>		50,000	

### OAR 340-041-8033 Table 31 Aquatic Life Water Quality Guidance Values for Toxic Pollutants

EPA	Ballistans	040	Fresh	water	Salt	water
No.	<del>Pollutant</del>	CAS Number	Acute	Chronic	Acute	Chronic
	Chlorinated Benzenes		<del>250</del>	<del>50</del>	<del>160</del>	<del>129</del>
	Chlorinated naphthalenes		1,600		<del>7.5</del>	
	Chloroalkyl Ethers		238,000			
<del>26</del>	Chloroform	<del>67663</del>	28,900	1,240		
<del>45</del>	Chlorophenol 2-	95578	4,380	2,000		
	Chlorophenol 4-	106489			<del>29,700</del>	
<del>52</del>	Methyl-4-chlorophenol 3-	<del>59507</del>	<del>30</del>			
<del>5a</del>	Chromium (III)	<del>1606583</del> <del>1</del>			10,300	
<del>109</del>	DDE 4,4'-	<del>72559</del>	<del>1,050</del>		14	
<del>110</del>	DDD-4,4'-	<del>72548</del>	0.06		<del>3.6</del>	
	Diazinon	<del>333415</del>	0.08	0.05		
	Dichlorobenzenes		1,120	<del>763</del>	1,970	
<del>29</del>	Dichloroethane 1,2-	107062	118,000	20,000	<del>113,00</del> <del>0</del>	
	Dichloroethylenes		11,600		<del>224,00</del> <del>0</del>	
<del>46</del>	Dichlorophenol 2,4-	120832	<del>2,020</del>	<del>365</del>		
<del>31</del>	Dichloropropane 1,2-	<del>78875</del>	23,000	5,700	10,300	3,040
<del>32</del>	Dichloropropene 1,3-	<del>542756</del>	6,060	<del>244</del>	<del>790</del>	
47	Dimethylphenol 2,4-	<del>105679</del>	<del>2,120</del>			
	Dinitrotoluene		<del>330</del>	<del>230</del>	<del>590</del>	<del>370</del>
<del>16</del>	Dioxin (2,3,7,8-TCDD)	<del>1746016</del>	0.01	38 pg/L		

### OAR 340-041-8033 Table 31 Aquatic Life Water Quality Guidance Values for Toxic Pollutants

EPA	Pollutant	CAS Number	Freshwater		Saltwater	
No.			Acute	Chronic	Acute	Chronic
<del>85</del>	Diphenylhydrazine 1,2-	<del>122667</del>	<del>270</del>			
33	Ethylbenzene	100414	32,000		430	
86	Fluoranthene	206440	3,980		40	<del>16</del>
	Haloethers		<del>360</del>	<del>122</del>		
	Halomethanes		11,000		12,000	6,400
<del>89</del>	Hexachlorobutadiene	<del>87683</del>	<del>90</del>	9.3 32		
90	Hexachlorocyclopenta diene	77474	7	<del>5.2</del>	7	
91	Hexachloroethane	67721	<del>980</del>	<del>540</del>	940	
93	Isophorone	<del>78591</del>	117,000		12,900	
94	Naphthalene	91203	<del>2,300</del>	<del>620</del>	<del>2,350</del>	
<del>95</del>	Nitrobenzene	98953	27,000		6,680	
	Nitrophenols		<del>230</del>	<del>150</del>	4,850	
<del>26 B</del>	Nitrosamines	3557691 4	5,850		3,300,0 00	
	Pentachlorinated ethanes	-	<del>7,240</del>	1,100	<del>390</del>	<del>281</del>
<del>54</del>	Phenol	108952	10,200	<del>2,560</del>	5,800	_
	Phthalate esters	_	940	3	2,944	3.4
	Polynuclear Aromatic Hydrocarbons	-	-	-	<del>300</del>	-
	Tetrachlorinated Ethanes	-	9,320	-	-	-
<del>37</del>	Tetrachloroethane 1,1,2,2-	<del>79345</del>	-	<del>2,400</del>	9,020	-

### OAR 340-041-8033 Table 31 Aquatic Life Water Quality Guidance Values for Toxic Pollutants

EPA	Pollutant	CAS Number	Fresh	water	Saltwater	
No.			Acute	Chronic	Acute	Chronic
	Tetrachloroethanes	_	9,320	ı	ı	-
<del>38</del>	Tetrachloroethylene	127184	<del>5,280</del>	<del>840</del>	<del>10,200</del>	<del>450</del>
	Tetrachlorophenol 2,3,5,6	-	-	-	-	440
<del>12</del>	<del>Thallium</del>	7440280	<del>1,400</del>	<del>40</del>	<del>2,130</del>	-
<del>39</del>	Toluene	108883	<del>17,500</del>	-	<del>6,300</del>	<del>5,000</del>
	Trichlorinated ethanes	_	<del>18,000</del>	_	-	-
41	Trichloroethane 1,1,1-	<del>71556</del>	-	_	31,200	-
42	Trichloroethane 1,1,2-	<del>79005</del>	-	<del>9,400</del>	-	_
43	Trichloroethylene	<del>79016</del>	45,000	<del>21,900</del>	<del>2,000</del>	-
<del>55</del>	Trichlorophenol 2,4,6-	<del>88062</del>	-	970	-	-

The following chemicals/compounds/classes are of concern due to the potential for toxic effects to aquatic organisms; however, no guidance values are designated. If these compounds are identified in the waste stream, then a review of the scientific literature may be appropriate for deriving guidance values.

- □ Polybrominated diphenyl ethers (PBDE)
- **□**-Polybrominated biphenyls (PBB)
- **□** Pharmaceuticals
- □ Personal care products
- **□**-Alkyl Phenols
- Other chemicals with Toxic effects

### Footnotes:

- A Values in Table 31 are applicable to all basins.
- B This number was assigned to the list of non-priority pollutants in National Recommended Water Quality Criteria: 2002 (EPA-822-R-02-047).



Effective April 18, 2014

### **Human Health Criteria Summary**

The concentration for each pollutant listed in Table 40 was derived to protect Oregonians from potential adverse health impacts associated with long-term exposure to toxic substances associated with consumption of fish, shellfish, and water. The "organism only" criteria are established to protect fish and shellfish consumption and apply to waters of the state designated for fishing. The "water + organism" criteria are established to protect the consumption of drinking water, fish, and shellfish, and apply where both fishing and domestic water supply (public and private) are designated uses. All criteria are expressed as micrograms per liter (µg/L), unless otherwise noted. Pollutants are listed in alphabetical order. Additional information includes the Chemical Abstract Service (CAS) number, whether the criterion is based on carcinogenic effects (can cause cancer in humans), and whether there is an aquatic life criterion for the pollutant (i.e. "y"= yes, "n" = no). All the human health criteria were calculated using a fish consumption rate of 175 grams per day unless otherwise noted. A fish consumption rate of 175 grams per day is approximately equal to 23 8-ounce fish meals per month. For pollutants categorized as carcinogens, values represent a cancer risk of one additional case of cancer in one million people (i.e. 10<sup>-6</sup>), unless otherwise noted. All metals criteria are for total metal concentration, unless otherwise noted. Italicized pollutants represent non-priority pollutants. The human health criteria revisions established by OAR 340-041-0033 and shown in Table 40 do not become applicable for purposes of ORS chapter 468B or the federal Clean Water Act until approved by EPA pursuant to 40 CFR 131.21 (4/27/2000).

# OAR 340-041-8033 Table 40 Human Health Water Quality Criteria for Toxic Pollutants

					Human Health Criteria for t Consumption of:			
No.	Pollutant	CAS Number	Carcinogen	Aquatic Life Criterion	Water + Organism (µg/L)	Organism Only (μg/L)		
1	Acenaphthene	83329	n	n	95	99		
2	Acrolein	107028	n	n	0.88	0.93		
3	Acrylonitrile	107131	у	n	0.018	0.025		
4	Aldrin	309002	у	у	0.0000050	0.0000050		
5	Anthracene	120127	n	n	2900	4000		
6	Antimony	7440360	n	n	5.1	64		
7	Arsenic (inorganic) <sup>A</sup>	7440382	у	у	2.1	2.1(freshwater) 1.0 (saltwater)		
A The arsenic criteria are expressed as total inorganic arsenic. The "organism only" freshwater criterion is based on a risk level of approximately 1 x 10 <sup>-5</sup> , and the "water + organism" criterion is based on a risk level of 1 x 10 <sup>-4</sup> .								
8	Asbestos <sup>B</sup>	1332214	у	n	7,000,000 fibers/L			
B The human health risks from asbestos are primarily from drinking water, therefore no "organism only" criterion was developed. The "water + organism" criterion is based on the Maximum Contaminant Level (MCL) established under the Safe Drinking Water Act.								
9	Barium <sup>c</sup>	7440393	n	n	1000			
<sup>c</sup> The human health criterion for barium is the same as originally published in the 1976 EPA Red Book which predates the 1980 methodology and did not utilize the fish ingestion BCF approach. This same criterion value was also published in the 1986 EPA Gold Book. Human health risks are primarily from drinking water, therefore no "organism only" criterion was developed. The "water + organism" criterion is based on the Maximum Contaminant Level (MCL) established under the Safe Drinking Water Act.								
10	Benzene	71432	у	n	0.44	1.4		
11	Benzidine	92875	у	n	0.000018	0.000020		
12	Benz(a)anthracene	56553	у	n	0.0013	0.0018		

					Human Health Criteria for the Consumption of:	
No.	Pollutant	CAS Number	Carcinogen	Aquatic Life Criterion	Water + Organism (µg/L)	Organism Only (μg/L)
13	Benzo(a)pyrene	50328	у	n	0.0013	0.0018
14	Benzo(b)fluoranthene 3,4	205992	у	n	0.0013	0.0018
15	Benzo(k)fluoranthene	207089	у	n	0.0013	0.0018
16	BHC Alpha	319846	У	n	0.00045	0.00049
17	BHC Beta	319857	у	n	0.0016	0.0017
18	BHC Gamma (Lindane)	58899	n	у	0.17	0.18
19	Bromoform	75252	у	n	3.3	14
20	Butylbenzyl Phthalate	85687	n	n	190	190
21	Carbon Tetrachloride	56235	у	n	0.10	0.16
22	Chlordane	57749	у	у	0.000081	0.000081
23	Chlorobenzene	108907	n	n	74	160
24	Chlorodibromomethane	124481	у	n	0.31	1.3
25	Chloroethyl Ether bis 2	111444	у	n	0.020	0.053
26	Chloroform	67663	n	n	260	1100
27	Chloroisopropyl Ether bis 2	108601	n	n	1200	6500
28	Chloromethyl ether, bis	542881	у	n	0.000024	0.000029
29	Chloronaphthalene 2	91587	n	n	150	160
30	Chlorophenol 2	95578	n	n	14	15

	Human Health Water Quality Criteria for Toxic Pollutants								
					Human Health ( Consum				
No.	Pollutant	CAS Number	Carcinogen	Aquatic Life Criterion	Water + Organism (µg/L)	Organism Only (µg/L)			
31	Chlorophenoxy Herbicide (2,4,5,-TP) <sup>D</sup>	93721	n	n	10				
<sup>D</sup> The Chlorophenoxy Herbicide (2,4,5,-TP) criterion is the same as originally published in the 1976 EPA Red Book which predates the 1980 methodology and did not utilize the fish ingestion BCF approach. This same criterion value was also published in the 1986 EPA Gold Book. Human health risks are primarily from drinking water, therefore no "organism only" criterion was developed. The "water + organism" criterion is based on the Maximum Contaminant Level (MCL) established under the Safe Drinking Water Act.									
32	Chlorophenoxy Herbicide (2,4-D) <sup>E</sup>	94757	n	n	100				
E The Chlorophenoxy Herbicide (2,4-D) criterion is the same as originally published in the 1976 EPA Red Book which predates the 1980 methodology and did not utilize the fish ingestion BCF approach. This same criterion value was also published in the 1986 EPA Gold Book. Human health risks are primarily from drinking water, therefore no "organism only" criterion was developed. The "water + organism" criterion is based on the Maximum Contaminant Level (MCL) established under the Safe Drinking Water Act.									
33	Chrysene	218019	у	n	0.0013	0.0018			
34	Copper <sup>F</sup>	7440508	n	у	1300				
	nan health risks from copper are prim r + organism" criterion is based on th								
35	Cyanide <sup>G</sup>	57125	n	у	130	130			

	"water + organism" criterion is based on the Maximum Contaminant Level (MCL) established under the Safe Drinking Water Act.								
35	Cyanide <sup>G</sup>	57125	n	у	130	130			
	<sup>G</sup> The cyanide criterion is expressed as total cyanide (CN)/L.								
36	DDD 4,4'	72548	у	n	0.000031	0.000031			
37	DDE 4,4'	72559	у	n	0.000022	0.000022			
38	DDT 4,4'	50293	у	у	0.000022	0.000022			
39	Dibenz(a,h)anthracene	53703	у	n	0.0013	0.0018			
40	Dichlorobenzene(m) 1,3	541731	n	n	80	96			

					Human Health Criteria for the Consumption of:	
No.	Pollutant	CAS Number	Carcinogen	Aquatic Life Criterion	Water + Organism (µg/L)	Organism Only (μg/L)
41	Dichlorobenzene(o) 1,2	95501	n	n	110	130
42	Dichlorobenzene(p) 1,4	106467	n	n	16	19
43	Dichlorobenzidine 3,3'	91941	у	n	0.0027	0.0028
44	Dichlorobromomethane	75274	у	n	0.42	1.7
45	Dichloroethane 1,2	107062	у	n	0.35	3.7
46	Dichloroethylene 1,1	75354	n	n	230	710
47	Dichloroethylene trans 1,2	156605	n	n	120	1000
48	Dichlorophenol 2,4	120832	n	n	23	29
49	Dichloropropane 1,2	78875	у	n	0.38	1.5
50	Dichloropropene 1,3	542756	у	n	0.30	2.1
51	Dieldrin	60571	у	у	0.0000053	0.0000054
52	Diethyl Phthalate	84662	n	n	3800	4400
53	Dimethyl Phthalate	131113	n	n	84000	110000
54	Dimethylphenol 2,4	105679	n	n	76	85
55	Di-n-butyl Phthalate	84742	n	n	400	450
56	Dinitrophenol 2,4	51285	n	n	62	530
57	Dinitrophenols	25550587	n	n	62	530
58	Dinitrotoluene 2,4	121142	у	n	0.084	0.34

					Human Health Criteria for the Consumption of:	
No.	Pollutant	CAS Number	Carcinogen	Aquatic Life Criterion	Water + Organism (µg/L)	Organism Only (μg/L)
59	Dioxin (2,3,7,8-TCDD)	1746016	у	n	0.00000000051	0.00000000051
60	Diphenylhydrazine 1,2	122667	у	n	0.014	0.020
61	Endosulfan Alpha	959988	n	У	8.5	8.9
62	Endosulfan Beta	33213659	n	У	8.5	8.9
63	Endosulfan Sulfate	1031078	n	n	8.5	8.9
64	Endrin	72208	n	у	0.024	0.024
65	Endrin Aldehyde	7421934	n	n	0.030	0.030
66	Ethylbenzene	100414	n	n	160	210
67	Ethylhexyl Phthalate bis 2	117817	у	n	0.20	0.22
68	Fluoranthene	206440	n	n	14	14
69	Fluorene	86737	n	n	390	530
70	Heptachlor	76448	у	у	0.0000079	0.0000079
71	Heptachlor Epoxide	1024573	у	у	0.0000039	0.0000039
72	Hexachlorobenzene	118741	у	n	0.000029	0.000029
73	Hexachlorobutadiene	87683	у	n	0.36	1.8
74	Hexachlorocyclo-hexane- Technical	608731	у	n	0.0014	0.0015
75	Hexachlorocyclopentadiene	77474	n	n	30	110
76	Hexachloroethane	67721	у	n	0.29	0.33

					Human Health Criteria for the Consumption of:	
No.	Pollutant	CAS Number	Carcinogen	Aquatic Life Criterion	Water + Organism (µg/L)	Organism Only (μg/L)
77	Indeno(1,2,3-cd)pyrene	193395	у	n	0.0013	0.0018
78	Isophorone	78591	у	n	27	96
79	Manganese <sup>H</sup>	7439965	n	n		100

<sup>&</sup>lt;sup>H</sup> The "fish consumption only" criterion for manganese applies only to salt water and is for total manganese. This EPA recommended criterion predates the 1980 human health methodology and does not utilize the fish ingestion BCF calculation method or a fish consumption rate.

80	Methoxychlor <sup>I</sup>	72435	n	у	100	
----	---------------------------	-------	---	---	-----	--

<sup>&</sup>lt;sup>1</sup> The human health criterion for methoxychlor is the same as originally published in the 1976 EPA Red Book which predates the 1980 methodology and did not utilize the fish ingestion BCF approach. This same criterion value was also published in the 1986 EPA Gold Book. Human health risks are primarily from drinking water, therefore no "organism only" criterion was developed. The "water + organism" criterion is based on the Maximum Contaminant Level (MCL) established under the Safe Drinking Water Act.

81	Methyl Bromide	74839	n	n	37	150
82	Methyl-4,6-dinitrophenol 2	534521	n	n	9.2	28
83	Methylene Chloride	75092	У	n	4.3	59
84	Methylmercury (mg/kg) <sup>J</sup>	22967926	n	n		0.040 mg/kg

<sup>&</sup>lt;sup>J</sup> This value is expressed as the fish tissue concentration of methylmercury. Contaminated fish and shellfish is the primary human route of exposure to methylmercury.

85	Nickel	7440020	n	У	140	170
86	Nitrates <sup>K</sup>	14797558	n	n	10000	

K The human health criterion for nitrates is the same as originally published in the 1976 EPA Red Book which predates the 1980 methodology and did not utilize the fish ingestion BCF approach. This same criterion value was also published in the 1986 EPA Gold Book. Human health risks are primarily from drinking water, therefore no "organism only" criterion was developed. The "water + organism" criterion is based on the Maximum Contaminant Level (MCL) established under the Safe Drinking Water Act.

					Human Health Criteria for the Consumption of:	
No.	Pollutant	CAS Number	Carcinogen	Aquatic Life Criterion	Water + Organism (µg/L)	Organism Only (μg/L)
87	Nitrobenzene	98953	n	n	14	69
88	Nitrosamines	35576911	У	n	0.00079	0.046
89	Nitrosodibutylamine, N	924163	У	n	0.0050	0.022
90	Nitrosodiethylamine, N	55185	У	n	0.00079	0.046
91	Nitrosodimethylamine, N	62759	У	n	0.00068	0.30
92	Nitrosodi-n-propylamine, N	621647	У	n	0.0046	0.051
93	Nitrosodiphenylamine, N	86306	у	n	0.55	0.60
94	Nitrosopyrrolidine, N	930552	у	n	0.016	3.4
95	Pentachlorobenzene	608935	n	n	0.15	0.15
96	Pentachlorophenol	87865	у	у	0.15	0.30
97	Phenol	108952	n	n	9400	86000
98	Polychlorinated Biphenyls (PCBs) <sup>L</sup>	NA	у	у	0.000064	0.0000064
	<sup>L</sup> This criterion a	pplies to total F	PCBs (e.g. determ	ined as Aroclo	rs or congeners).	
99	Pyrene	129000	n	n	290	400
100	Selenium	7782492	n	у	120	420
101	Tetrachlorobenzene, 1,2,4,5-	95943	n	n	0.11	0.11
102	Tetrachloroethane 1,1,2,2	79345	у	n	0.12	0.40
103	Tetrachloroethylene	127184	У	n	0.24	0.33

					Human Health Criteria for the Consumption of:		
No.	Pollutant	CAS Number	Carcinogen	Aquatic Life Criterion	Water + Organism (µg/L)	Organism Only (μg/L)	
104	Thallium	7440280	n	n	0.043	0.047	
105	Toluene	108883	n	n	720	1500	
106	Toxaphene	8001352	у	у	0.000028	0.000028	
107	Trichlorobenzene 1,2,4	120821	n	n	6.4	7.0	
108	Trichloroethane 1,1,2	79005	у	n	0.44	1.6	
109	Trichloroethylene	79016	у	n	1.4	3.0	
110	Trichlorophenol 2,4,6	88062	у	n	0.23	0.24	
111	Trichlorophenol, 2, 4, 5-	95954	n	n	330	360	
112	Vinyl Chloride	75014	у	n	0.023	0.24	
113	Zinc	7440666	n	у	2100	2600	

### **Draft Rules – Edits Included**

### Division 41 WATER QUALITY STANDARDS: BENEFICIAL USES, POLICIES, AND CRITERIA FOR OREGON

### 340-041-0033 Toxic Substances

- (1) Toxic Substances Narrative. Toxic substances may not be introduced above natural background levels in waters of the state in amounts, concentrations, or combinations that may be harmful, may chemically change to harmful forms in the environment, or may accumulate in sediments or bioaccumulate in aquatic life or wildlife to levels that adversely affect public health, safety, or welfare or aquatic life, wildlife or other designated beneficial uses.
- (2) Aquatic Life Numeric Criteria. Levels of toxic substances in waters of the state may not exceed the applicable aquatic life criteria as defined in Table 30 under OAR 340-041-8033.
- (3) Human Health Numeric Criteria. The criteria for waters of the state listed in Table 40 under OAR 340-041-8033 are established to protect Oregonians from potential adverse health effects associated with long-term exposure to toxic substances associated with consumption of fish, shellfish and water.
- (4) To establish permit or other regulatory limits for toxic substances without criteria in Table 30 under OAR 340-041-8033 or Table 40 under 340-041-8033, DEQ may use public health advisories and published scientific literature. DEQ may also require or conduct bio-assessment studies to monitor the toxicity to aquatic life of complex effluents, other suspected discharges or chemical substances without numeric criteria.
- (5) Establishing Site-Specific Background Pollutant Criteria: This provision is a performance- based water quality standard that results in site-specific human health water quality criteria under the conditions and procedures specified in this rule section. It addresses existing permitted discharges of a pollutant removed from the same body of water. For waterbodies where a discharge does not increase the pollutant's mass and does not increase the pollutant concentration by more than 3 percent, and where the water body meets a pollutant concentration associated with a risk level of 1 x 10-4, DEQ concludes that the pollutant concentration continues to protect human health.
- (a) Definitions: As used in this section:

- (A) "Background pollutant concentration" means the ambient water body concentration immediately upstream of the discharge, regardless of whether those pollutants are natural or result from upstream human activity.
- (B) An "intake pollutant" is the amount of a pollutant present in waters of the state (including groundwater) as provided in subsection (C), below, at the time it is withdrawn from such waters by the discharger or other facility supplying the discharger with intake water.
- (C) "Same body of water": An intake pollutant is considered to be from the "same body of water" as the discharge if DEQ finds that the intake pollutant would have reached the vicinity of the outfall point in the receiving water within a reasonable period had the permittee not removed it. To make this finding, DEQ requires information showing that:
- (i) The background concentration of the pollutant in the receiving water (excluding any amount of the pollutant in the facility's discharge) is similar to that in the intake water; and.
- (ii) There is a direct hydrological connection between the intake and discharge points.
- (I) DEQ may also consider other site-specific factors relevant to the transport and fate of the pollutant to make the finding in a particular case that a pollutant would or would not have reached the vicinity of the outfall point in the receiving water within a reasonable period had the permittee not removed it.
- (II) An intake pollutant from groundwater may be considered to be from the "same body of water" if DEQ determines that the pollutant would have reached the vicinity of the outfall point in the receiving water within a reasonable period had the permittee not removed it. A pollutant is not from the same body of water if the groundwater contains the pollutant partially or entirely due to past or present human activity, such as industrial, commercial, or municipal operations, disposal actions, or treatment processes.
- (iii) Water quality characteristics (e.g., temperature, pH, hardness) are similar in the intake and receiving waters.
- (b) Applicability
- (A) DEQ may establish site-specific criteria under this rule section only for carcinogenic pollutants.
- (B) Site-specific criteria established under this rule section apply in the vicinity of the discharge for purposes of establishing permit limits for the specified permittee.
- (C) The underlying waterbody criteria continue to apply for all other Clean Water Act programs.

- (D) The site-specific background pollutant criterion will be effective when DEQ issues the permit for the specified permittee.
- (E) DEQ will reevaluate any site-specific criteria developed under this procedure upon permit renewal.
- (c) DEQ may establish a site-specific background pollutant criterion when all of the following conditions are met:
- (A) The discharger has a currently effective NPDES permit;
- (B) The mass of the pollutant discharged to the receiving waterbody does not exceed the mass of the intake pollutant from the same body of water, as defined in section (5)(a)(C) above, and therefore does not increase the total mass load of the pollutant in the receiving water body;
- (C) DEQ has not assigned the discharger a TMDL wasteload allocation for the pollutant in question;
- (D) The permittee uses any feasible pollutant reduction measures available and known to minimize the pollutant concentration in their discharge;
- (E) The pollutant discharge has not been chemically or physically altered in a manner that causes adverse water quality impacts that would not occur if the intake pollutants were left in-stream; and,
- (F) The timing and location of the pollutant discharge would not cause adverse water quality impacts that would not occur if the intake pollutant were left in-stream.
- (d) The site-specific background pollutant criterion must be the most conservative of the following four values. Section (5)(e) of this rule describes the procedures for deriving these values.
- (A) The projected in-stream pollutant concentration resulting from the current discharge concentration and any feasible pollutant reduction measures under (c)(D) above, after mixing with the receiving stream.
- (B) The projected in-stream pollutant concentration resulting from the portion of the current discharge concentration associated with the intake pollutant mass after mixing with the receiving stream. This analysis ensures that there will be no increase in the mass of the intake pollutant in the receiving water body as required by condition (c)(B) above.
- (C) The projected in-stream pollutant concentration associated with a 3 percent increase above the background pollutant concentration as calculated:

- (i) For the main stem Willamette and Columbia Rivers, using 25 percent of the harmonic mean flow of the waterbody.
- (ii) For all other waters, using 100 percent of the harmonic mean flow or similar critical flow value of the waterbody.
- (D) A criterion concentration value representing a human health risk level of 1 x 10-4. DEQ calculates this value using EPA's human health criteria derivation equation for carcinogens (EPA 2000), a risk level of 1 x 10-4, and the same values for the remaining calculation variables that were used to derive the underlying human health criterion.
- (e) Procedure to derive a site-specific human health water quality criterion to address a background pollutant:
- (A) DEQ will develop a flow-weighted characterization of the relevant flows and pollutant concentrations of the receiving waterbody, effluent and all facility intake pollutant sources to determine the fate and transport of the pollutant mass.
- (i) The pollutant mass in the effluent discharged to a receiving waterbody may not exceed the mass of the intake pollutant from the same body of water.
- (ii) Where a facility discharges intake pollutants from multiple sources that originate from the receiving waterbody and from other waterbodies, DEQ will calculate the flow-weighted amount of each source of the pollutant in the characterization.
- (iii) Where a municipal water supply system provides intake water for a facility and the supplier provides treatment of the raw water that removes an intake water pollutant, the concentration and mass of the intake water pollutant must be determined at the point where the water enters the water supplier's distribution system.
- (B) Using the flow weighted characterization developed in section (5)(e)(A), DEQ will calculate the in-stream pollutant concentration following mixing of the discharge into the receiving water. DEQ will use the resultant concentration to determine the conditions in section (5)(d)(A) and (B).
- (C) Using the flow-weighted characterization, DEQ will calculate the in-stream pollutant concentration based on an increase of 3 percent above background pollutant concentration. DEQ will use the resultant concentration to determine the condition in Section (5)(d)(C).
- (i) For the main stem Willamette and Columbia Rivers, DEQ will use 25 percent of the harmonic mean flow of the waterbody.
- (ii) For all other waters, DEQ will use 100 percent of the harmonic mean flow or similar critical flow value of the waterbody.

- (D) DEQ will select the most conservative of the following values as the site-specific water quality criterion.
- (i) The projected in-stream pollutant concentration described in section (5)(e)(B);
- (ii) The in-stream pollutant concentration based on an increase of 3 percent above background described in section (5)(e)(C); or
- (iii) A water quality criterion based on a risk level of 1 x 10-4.
- (f) Calculation of water quality based effluent limits based on a site-specific background pollutant criterion:
- (A) For discharges to receiving waters with a site-specific background pollutant criterion, DEQ will use the site-specific criterion in the calculation of a numeric water quality based effluent limit.
- (B) DEQ will compare the calculated water quality based effluent limits to any applicable aquatic toxicity or technology based effluent limits and select the most conservative for inclusion in the permit conditions.
- (g) In addition to the water quality based effluent limits described in section (5)(f), DEQ will calculate a mass-based limit where necessary to ensure that the condition described in section (5)(c)(B) is met. Where mass-based limits are included, the permit will specify how DEQ will assess compliance with mass-based effluent limitations.
- (h) The permit shall include a provision requiring DEQ to consider the re-opening of the permit and re-evaluation of the site-specific background pollutant criterion if new information shows the discharger no longer meets the conditions described in subsections (5)(c) and (e).
- (i) Public Notification Requirements.
- (A) If DEQ proposes to grant a site-specific background pollutant criterion, it must provide public notice of the proposal and hold a public hearing. The public notice may be included in the public notification of a draft NPDES permit or other draft regulatory decision that would rely on the criterion and will also be published on DEQ's water quality standards website;
- (B) DEQ will publish a list of all site-specific background pollutant criteria approved according to this rule. DEQ will add the criterion to this list within 30 days of its effective date. The list will identify the:
- (i) Permittee;
- (ii) Site-specific background pollutant criterion and the associated risk level;

- (iii) Waterbody to which the criterion applies;
- (iv) Allowable pollutant effluent limit; and,
- (v) How to obtain additional information about the criterion.
- (6) Arsenic Reduction Policy: The inorganic arsenic criterion for the protection of human health from the combined consumption of organisms and drinking water is 2.1 micrograms per liter. While this criterion is protective of human health and more stringent than the federal maximum contaminant level (MCL) for arsenic in drinking water, which is 10 micrograms per liter, it is based on a higher risk level than EQC used to establish other human health criteria. This higher risk level recognizes that much of the risk is due to naturally high levels of inorganic arsenic in Oregon's waterbodies. In order to maintain the lowest human health risk from inorganic arsenic in drinking water, EQC determined that it is appropriate to adopt the following policy to limit the human contribution to that risk.
- (a) It is EQC policy to reduce the addition of inorganic arsenic from new or existing anthropogenic sources to waters of the state within a surface water drinking water protection area to the maximum amount feasible. The requirements of this rule section (OAR 340-041-0033(6)) apply to sources that discharge to surface waters of the state with an ambient inorganic arsenic concentration equal to or lower than the applicable numeric inorganic arsenic criteria for the protection of human health.
- (b) Definitions. As used in this section:
- (A) "Add inorganic arsenic" means to discharge a net mass of inorganic arsenic from a point source (the mass of inorganic arsenic discharged minus the mass of inorganic arsenic taken into the facility from a surface water source).
- (B) A "surface water drinking water protection area," means an area delineated as such by DEQ under the source water assessment program of the federal Safe Drinking Water Act, 42 U.S.C. § 300j 13. DEQ delineates these areas to protect public or community drinking water supplies that use surface water sources. These delineations are on DEQ's drinking water program Web page.
- (C) "Potential to significantly increase inorganic arsenic concentrations in the public drinking water supply source water" means:
- (i) A discharge will increase the concentration of inorganic arsenic in the receiving water by 10 percent or more after mixing with the harmonic mean flow of the receiving water; or
- (ii) As an alternative, if sufficient data are available, the discharge will increase the concentration of inorganic arsenic in the surface water intake water of a public water system by 0.021 micrograms per liter or more based on a mass balance calculation.

- (c) Following the effective date of this rule, applications for an individual NPDES permit or permit renewal received from industrial dischargers located in a surface water drinking water protection area and identified by DEQ as likely to add inorganic arsenic to the receiving water must include sufficient data to enable DEQ to determine whether:
- (A) The discharge adds inorganic arsenic; and,
- (B) The discharge has the potential to significantly increase inorganic arsenic concentrations in the public drinking water supply source water.
- (d) Where DEQ determines that both conditions in subsection (c) of this section (6) are true, the industrial discharger must develop an inorganic arsenic reduction plan and propose all feasible measures to reduce its inorganic arsenic loading to the receiving water. The proposed plan, including proposed measures, monitoring and reporting requirements, and a schedule for those actions, will be described in the fact sheet and incorporated into the source's NPDES permit after public comment and DEQ review and approval. In developing the plan, the source must:
- (A) Identify how much it can minimize its inorganic arsenic discharge through pollution prevention measures, process changes, wastewater treatment, alternative water supply for groundwater users, or other possible pollution prevention and control measures;
- (B) Evaluate the costs, feasibility and environmental impacts of the potential inorganic arsenic reduction and control measures;
- (C) Estimate the predicted reduction in inorganic arsenic and the reduced human health risk expected to result from the control measures;
- (D) Propose specific inorganic arsenic reduction or control measures, if feasible, and an implementation schedule; and,
- (E) Propose monitoring and reporting requirements to document progress in plan implementation and the inorganic arsenic load reductions.
- (e) In order to implement this section, DEQ will develop the following information and guidance within 120 days of the effective date of this rule and periodically update it as warranted by new information:
- (A) A list of industrial sources or source categories, including industrial stormwater and sources covered by general permits likely to add inorganic arsenic to surface waters of the state. For industrial sources or source categories permitted under a general permit that have been identified by DEQ as likely sources of inorganic arsenic, DEQ will evaluate options for reducing inorganic arsenic during permit renewal or evaluation of Stormwater Pollution Control Plans.
- (B) Quantitation limits for monitoring inorganic arsenic concentrations.

- (C) Information and guidance to assist sources in estimating, according to subsection (d)(C) of this section, the reduced human health risk expected to result from inorganic arsenic control measures based on the most current EPA risk assessment.
- (f) It is the policy of EQC that landowners engaged in agricultural or development practices on land where pesticides, fertilizers, or soil amendments containing arsenic are currently being or have previously been applied, implement conservation practices to minimize the erosion and runoff of inorganic arsenic to waters of the state or to a location where such material could readily migrate into waters of the state.

[NOTE: Tables 30 and 40 are found under OAR 340-041-8033]

**Statutory/Other Authority:** ORS 468.020, 468B.030, 468B.035 & 468B.048 **Statutes/Other Implemented:** ORS 468B.030, 468B.035 & 468B.048

**History:** 

DEQ 11-2016, f. & cert. ef. 11-2-16

DEQ 1-2015, f. & cert. ef. 1-7-15

DEQ 17-2013, f. 12-23-13, cert. ef. 4-18-14

DEQ 10-2011, f. & cert. ef. 7-13-11

DEQ 8-2011, f. & cert. ef. 6-30-11

DEQ 17-2010, f. & cert. ef. 12-21-10

DEQ 3-2004, f. & cert. ef. 5-28-04

DEQ 17-2003, f. & cert. ef. 12-9-03

### Division 41 WATER QUALITY STANDARDS: BENEFICIAL USES, POLICIES, AND CRITERIA FOR OREGON

### 340-041-8033 Division 41 Tables and Figures

- (1) Table 30: Aquatic Life Water Quality Criteria for Toxic Pollutants. This table, referenced in OAR 340-041-0033, contains information about the applicability and content of the criteria contained in the table.
- (2) Table 40: Human Health Water Quality Criteria for Toxic Pollutants. This table, referenced in OAR 340-041-0033, contains information about the applicability and content of the criteria contained in the table.

[ED. NOTE: To view attachments referenced in rule text, click here for PDF copy.] **Statutory/Other Authority**: ORS 468.020, 468B.030, 468B.035 & 468B.048Statutes/Other **Implemented**: ORS 468B.030, 468B.035 & 468B.048 **History**:

DEQ 13-2019, amend filed 05/16/2019, effective 05/16/2019

DEQ 13-2017, minor correction filed 10/30/2017, effective 10/30/2017

DEQ 11-2016, f. & cert. ef. 11-2-16

DEQ 1-2015, f. & cert. ef. 1-7-15



The concentration for each compound listed in Table 30 is a criterion established for waters of the state in order to protect aquatic life. The aquatic life criteria apply to waterbodies where the protection of fish and aquatic life is a designated use. All values are expressed as micrograms per liter ( $\mu$ g/L). Compounds are listed in alphabetical order with the corresponding information: the Chemical Abstract Service (CAS) number, whether there is a human health criterion for the pollutant (i.e. "y"= yes, "n" = no), and the associated aquatic life freshwater and saltwater acute and chronic criteria. *Italicized* pollutants are not identified as priority pollutants by EPA. Dashes in the table column indicate that there is no aquatic life criterion.

Unless otherwise noted in the table below, the acute criterion is the Criterion Maximum Concentration (CMC) applied as a one-hour average concentration, and the chronic criterion is the Criterion Continuous Concentration (CCC) applied as a 96-hour (4 days) average concentration. The CMC and CCC criteria may not be exceeded more than once every three years. Footnote A, associated with eleven pesticide pollutants in Table 30, describes the exception to the frequency and duration of the toxics criteria stated in this paragraph.

	•			Fresh (µg		Saltwater (µg/L)			
No.	Pollutant	CAS Number	Human Health Criterion	Acute Criterion (CMC)	Chronic Criterion (CCC)	Acute Criterion (CMC)	Chronic Criterion (CCC)		
1	Acrolein	107028	y	3.0	3.0				
NOT	<mark>E: These acrolein c</mark>	riteria are n	ot applicable	e for Clean Wat	t <mark>er Act purpose</mark>	es until approve	ed by EPA.		
<mark>2</mark>	Aldrin	309002	у	3 A		1.3 <sup>A</sup>			
<sup>A</sup> See	<sup>A</sup> See expanded endnote A at bottom of Table 30 for alternate frequency and duration of this criterion.								
3	Alkalinity		n		20,000 <sup>B</sup>				

				Fresh (µg			twater ug/L)
No.	Pollutant	CAS Number	Human Health Criterion	Acute Criterion (CMC)	Chronic Criterion (CCC)	Acute Criterion (CMC)	Chronic Criterion (CCC)
	B Criterion shown is	s the minimum	(i.e. CCC in w	ater may not be be	elow this value in (	order to protect aq	uatic life).
4	<u>Aluminum</u>	7429905	n	See O, P	See O, P		<mark></mark>

O The freshwater criterion for aluminum is a function of the pH, dissolved organic carbon, and total hardness in the water column. Acute (CMC) and chronic (CCC) freshwater aluminum criteria values shall be calculated using the 2018 Aluminum Criteria Calculator (Aluminum Criteria Calculator V.2.0.xlsx), or a calculator in R or other software package using the same 1985 Guidelines calculation approach and underlying model equations as in the Aluminum Criteria Calculator V.2.0.xlsx, as defined in EPA's Final Aquatic Life Ambient Water Quality Criteria for Aluminum (EPA 822-R-18-001). See also endnote O for procedures and information.

P Oregon will use analytical methods that measure the bioavailable fraction of aluminum unless total recoverable aluminum measurements are required by Federal regulations.

Note: These aluminum criteria are not applicable for CWA purposes until approved by EPA and, if necessary, EPA withdraws the federal criteria for Oregon. The currently applicable federally promulgated criteria may be found at 40 CFR 131.47.

<mark>5</mark>	Ammonia	7664417	n	The ammonia criteria are pH and temperature dependent — See ammonia criteria Tables 30(a)-(c) at end of Table 30. <sup>M</sup>	The ammonia criteria are pH, temperature and salinity dependent. Values for saltwater criteria (total ammonia) can be calculated from the tables specified in Ambient Water Quality Criteria for Ammonia (Saltwater)—1989 (EPA 440/5-88-004) See DEQ's calculator for calculating saltwater ammonia criteria at:

				Fresh (µg			water g/L)
No.	Pollutant	CAS Number	Human Health Criterion	Acute Criterion (CMC)	Chronic Criterion (CCC)	Acute Criterion (CMC)	Chronic Criterion (CCC)
						http://www.de q/standards/to	eq.state.or.us/w oxics.htm

M The acute criteria in Table 30(a) apply in waterbodies where salmonids are a designated use in OAR 340-041-0101 through OAR 340-041-0340. The acute criteria in Table 30(b) apply in waterbodies where salmonids are not a designated use. The chronic criteria in Table 30(c) apply where fish and aquatic life is a designated use. It is not necessary to account for the presence or absence of salmonids or the presence of any early life stage of fish for the chronic criteria. Refer to DEQ's beneficial use website at: http://www.deq.state.or.us/wq/standards/uses.htm for additional information on salmonid beneficial use designations, including tables and maps.

						36 C, D	
	terion is expressed terion is applied as						
<mark>7</mark>	BHC Gamma (Lindane)	58899	у	0.95	0.08 <sup>A</sup>	0.16 <sup>A</sup>	
<sup>A</sup> Sec	e expanded endnot	e A at botto	m of Table 3	30 for alternate	frequency and	duration of this	s criterion.
8	Cadmium	7440439	n	See C, F	See C, F	33 <sup>c</sup>	<mark>7.9 <sup>c</sup></mark>

<sup>&</sup>lt;sup>c</sup> Criterion is expressed in terms of "dissolved" concentrations in the water column.

NOTE: The federally promulgated freshwater acute cadmium criterion in 40 CFR 131.46 remains applicable for CWA purposes until EPA approves the freshwater acute cadmium criterion specified in Table 30 Endnotes C and F. The following saltwater cadmium criteria remain applicable until EPA approves the revised criteria in Table 30: the cadmium acute (CMC) is 40 µg/L and the chronic (CCC) is 8.8 µg/L, both expressed in terms of "dissolved" concentrations in the water column.

|--|

NOTE: These carbaryl criteria are not applicable for Clean Water Act purposes until approved by EPA.

F The freshwater criterion for this metal is expressed as a function of hardness (mg/L) in the water column. To calculate the criterion, use the formula under expanded endnote F at bottom of Table 30.

				Fresh ( <i>µ</i> g			water g/L)
No.	Pollutant	CAS Number	Human Health Criterion	Acute Criterion (CMC)	Chronic Criterion (CCC)	Acute Criterion (CMC)	Chronic Criterion (CCC)
<mark>10</mark>	Chlordane	57749	у	2.4 <sup>A</sup>	0.0043 A	0.09 A	0.004 <sup>A</sup>
<sup>A</sup> See	e expanded endnot	e A at botto	m of Table 3	30 for alternate	frequency and	duration of this	s criterion.
<mark>11</mark>	Chloride	1688700 6	n	860,000	230,000		
<mark>12</mark>	Chlorine	7782505	n	19	11	13	7.5
<mark>13</mark>	Chlorpyrifos	2921882	n	0.083	0.041	0.011	0.0056
<mark>14</mark>	Chromium III	1606583 1	n	See C, F	See C, F		
F The To ca	erion is expressed e freshwater criterion alculate the criterion	on for this m n, use formu	etal is expre	essed as a fund panded endnot	etion of hardnes e F at bottom o	ss (mg/L) in the of Table 30.	
<mark>15</mark>	Chromium VI	1854029 9	n	16 <sup>c</sup>	11 <sup>C</sup>	1100 <sup>c</sup>	<sub>50</sub> <b>c</b>
<sup>c</sup> Crit	erion is expressed	in terms of '	'dissolved" c	concentrations i	in the water co	lumn.	
<mark>16</mark>	Copper	7440508	у	See C, N	See C, N	4.8 <sup>c</sup>	3.1 °C
N The	erion is expressed freshwater criterio emperature in the vote N at the bottom	n for coppe water colum	r is a functio n. To calcula	n of the concer ate the criterion	ntration of ions, , use the Biotic	alkalinity, orga Ligand Model	referenced in

<sup>&</sup>lt;sup>N</sup> The freshwater criterion for copper is a function of the concentration of ions, alkalinity, organic carbon, pH and temperature in the water column. To calculate the criterion, use the Biotic Ligand Model referenced in endnote N at the bottom of Table 30. The acute copper criterion (CMC) is applied as a one-hour average concentration. The chronic criterion (CCC) is applied as a 96-hour (4 days) average concentration. See endnote N also for procedures and information.

<mark>17</mark>	Cyanide	57125	у	22 <sup>J</sup>	5.2 <sup>J</sup>	1 <sup>J</sup>	1 <sup>J</sup>
		<sup>J</sup> This crit	erion is exp	ressed as µg fr	ee cyanide (Cl	V)/L.	

				Fresh (µg	water g/L)		water g/L)
No.	Pollutant	CAS Number	Human Health Criterion	Acute Criterion (CMC)	Chronic Criterion (CCC)	Acute Criterion (CMC)	Chronic Criterion (CCC)
<mark>18</mark>	DDT 4,4'	50293	у	1.1 <sup>A, G</sup>	0.001 <sup>A, G</sup>	0.13 <sup>A, G</sup>	0.001 <sup>A, G</sup>
<sup>G</sup> This	e expanded endnot s criterion applies to ld not exceed this v	DDT and i					
<mark>19</mark>	Demeton	8065483	n		0.1		0.1
<mark>20</mark>	<u>Diazinon</u>	333415	<mark>n</mark>	<mark>0.17</mark>	<mark>0.17</mark>	0.82	0.82
NOT	E: These diazinon d	criteria are n	ot applicabl	e for Clean Wa	ter Act purpos	es until approve	ed by EPA.
<mark>21</mark>	Dieldrin	60571	у	0.24	0.056	0.71 <sup>A</sup>	0.0019 <sup>A</sup>
<sup>A</sup> See	e expanded endnot	e A at botto	m of Table 3	30 for alternate	frequency and	duration of this	s criterion.
<mark>22</mark>	Endosulfan	115297	n	0.22 <sup>A, H</sup>	0.056 <sup>A, H</sup>	0.034 <sup>A, H</sup>	0.0087 <sup>A, H</sup>
<sup>H</sup> This	e expanded endnot s value is based on 5-80-046) and shou	the criterior	n published i	in Ambient Wat	ter Quality Crite	eria for Endosu	
<mark>23</mark>	Endosulfan Alpha	959988	у	0.22 A	0.056 <sup>A</sup>	0.034 <sup>A</sup>	0.0087 <sup>A</sup>
<sup>A</sup> See	e expanded endnot	e A at botto	m of Table 3	30 for alternate	frequency and	duration of this	s criterion.
<mark>24</mark>	Endosulfan Beta	3321365 9	у	0.22 <sup>A</sup>	0.056 <sup>A</sup>	0.034 <sup>A</sup>	0.0087 <sup>A</sup>
<sup>A</sup> See	e expanded endnot	e A at botto	m of Table 3	30 for alternate	frequency and	duration of this	s criterion.
<mark>25</mark>	Endrin	72208	у	0.086	0.036	0.037 A	0.0023 A
<sup>A</sup> Sec	e expanded endnot	e A at botto	m of Table 3	30 for alternate	frequency and	duration of this	s criterion.
<mark>26</mark>	Guthion	86500	n		0.01		0.01
<mark>27</mark>	Heptachlor	76448	у	0.52 A	0.0038 A	0.053 A	0.0036 A
<sup>A</sup> See	e expanded endnot	e A at botto	m of Table 3	30 for alternate	frequency and	duration of this	s criterion.

	-			_			
				Fresh ( <i>µ</i> g			twater ug/L)
No.	Pollutant	CAS Number	Human Health Criterion	Acute Criterion (CMC)	Chronic Criterion (CCC)	Acute Criterion (CMC)	Chronic Criterion (CCC)
<mark>28</mark>	Heptachlor Epoxide	1024573	у	0.52 <sup>A</sup>	0.0038 <sup>A</sup>	0.053 A	0.0036 <sup>A</sup>
<sup>A</sup> See	e expanded endnot	e A at botto	m of Table 3	30 for alternate	frequency and	duration of this	s criterion.
<mark>29</mark>	Iron (total)	7439896	n		1000		
<mark>30</mark>	Lead	7439921	n	See C, F	See C, F	210 <sup>c</sup>	8.1 <sup>c</sup>
F The	erion is expressed freshwater criterionalculate the criterion	n for this me	etal is expres	ssed as a funct	ion of hardnes	s (mg/L) in the	water column.
<mark>31</mark>	Malathion	121755	n		0.1		0.1
<mark>32</mark>	Mercury (total)	7439976	n	2.4	0.012	2.1	0.025
<mark>33</mark>	Methoxychlor	72435	у		0.03		0.03
<mark>34</mark>	Mirex	2385855	n		0.001		0.001
<mark>35</mark>	Nickel	7440020	у	See C, F	See C, F	74 <sup>c</sup>	8.2 <sup>c</sup>
F The	erion is expressed e freshwater criterio alculate the criterion	on for this m	etal is expre	ssed as a func	tion of hardnes	ss (mg/L) in the	water column.
<mark>36</mark>	Parathion	56382	n	0.065	0.013		
<mark>37</mark>	Pentachlorophe nol	87865	у	See I	See I	13	7.9
	shwater aquatic life llows: CMC=(exp(1	•	•	•		ction of pH, an	d are calculated
38	Phosphorus Elemental	7723140	n		-	1	0.1

				Fresh ( <i>µ</i> g			twater <i>ıg/L)</i>
No.	Pollutant	CAS Number	Human Health Criterion	Acute Criterion (CMC)	Chronic Criterion (CCC)	Acute Criterion (CMC)	Chronic Criterion (CCC)
<mark>39</mark>	Polychlorinated Biphenyls (PCBs)	NA	У	2 <sup>K</sup>	0.014 <sup>K</sup>	10 <sup>K</sup>	0.03 <sup>K</sup>
K Thi	s criterion applies t	o total PCBs	(e.g. deterr	nined as Arock	ors or congene	rs)	
<mark>40</mark>	Selenium	7782492	у	See C , L	4.6 <sup>c</sup>	290 <sup>c</sup>	71 <sup>C</sup>
L The	erion is expressed CMC=(1/[(f1/CMC) ded as selenite and ectively. See expan	31)+(f2/CMC selenate, re	2)]µg/L) * C spectively, a	F where f1 and and CMC1 and	f2 are the frac CMC2 are 185	tions of total se 5.9 µg/L and 12	
<mark>41</mark>	Silver	7440224	n	See C , F	0.10 <sup>c</sup>	1.9 <sup>c</sup>	
F The	erion is expressed e freshwater acute nn. To calculate the	criterion for	this metal is	expressed as	a function of h	ardness (mg/L)	
<mark>42</mark>	Sulfide	7783064	n		2		2

<mark>42</mark>	Sulfide Hydrogen	7783064	n		2		2
43	Sulfide Toxaphene	8001352	у	0.73	0.0002	0.21	0.0002
<mark>44</mark>	Tributyltin (TBT)	688733	n	0.46	0.072	0.42	0.0074

NOTE: The freshwater chronic and saltwater acute and chronic criteria for tributyltin are not applicable for Clean Water Act purposes until the revised Table 30 values are approved by EPA. The freshwater acute criterion is not changing and is applicable. The following tributyltin criteria remain applicable until EPA approves the revised criteria shown in Table 30: the freshwater chronic (CCC) is 0.063 µg/L, the saltwater acute (CMC) is 0.37 µg/L, and the saltwater chronic (CCC) is 0.01 µg/L.

|--|

<sup>&</sup>lt;sup>c</sup> Criterion is expressed in terms of "dissolved" concentrations in the water column.

F The freshwater criterion for this metal is expressed as a function of hardness (mg/L) in the water column. To calculate the criterion, use formula under expanded endnote F at bottom of Table 30.

### Expanded Endnotes A, F, N, O

### **Endnote A: Alternate Frequency and Duration for Certain Pesticides**

This criterion is based on EPA recommendations issued in 1980 that were derived using guidelines that differed from EPA's 1985 Guidelines which update minimum data requirements and derivation procedures. The CMC may not be exceeded at any time and the CCC may not be exceeded based on a 24-hour average. The CMC may be applied using a one hour averaging period not to be exceeded more than once every three years, if the CMC values given in Table 30 are divided by 2 to obtain a value that is more comparable to a CMC derived using the 1985 Guidelines.

### Endnote F: Equations for Hardness-Dependent Freshwater Metals Criteria and Conversion Factor Table

The freshwater criterion for this metal is expressed as dissolved with two significant figures, and is a function of hardness (mg/L) in the water column. Criteria values based on hardness are calculated using the following formulas (CMC refers to the acute criterion; CCC refers to the chronic criterion):

**CMC** = 
$$(exp(m_A*[ln(hardness)] + b_A))*CF$$

**CCC** = 
$$(exp(m_C*[ln(hardness)] + b_C))*CF$$

"CF" is the conversion factor used for converting a metal criterion expressed as the total recoverable fraction in the water column to a criterion expressed as the dissolved fraction in the water column.

Values for Calculating Hardness-Dependent Metals Criteria											
Chemical m <sub>A</sub> b <sub>A</sub> m <sub>C</sub> b <sub>C</sub>											
Cadmium	<mark>0.9789</mark>	<mark>-3.866</mark>	0.7409	-4.719							
Chromium III	0.8190	3.7256	0.8190	0.6848							
Lead	1.273	-1.460	1.273	-4.705							
Nickel	0.8460	2.255	0.8460	0.0584							
Silver	1.72	-6.59									
Zinc	0.8473	0.884	0.8473	0.884							

The conversion factors (CF) below must be used in the equations above for the hardness-dependent metals in order to convert total recoverable metals criteria to dissolved metals criteria. For metals that are not hardness-dependent (i.e. arsenic, chromium VI, selenium, and silver (chronic)), or are saltwater criteria, the criterion value associated with the metal in Table 30 already reflects a dissolved criterion based on its conversion factor below.

C	Conversion Factor (CI	Table for Dissolved	d Metals			
Chemical	Fresh	water	Saltwater			
Cileillicai	Acute	Chronic	Acute	Chronic		
Arsenic	1.000	1.000	1.000	1.000		
Cadmium	1.136672-[(ln hardness)(0.041838)]	1.101672-[(In hardness)(0.041838)]	0.994	0.994		
Chromium III	0.316	0.860				
Chromium VI	0.982	0.962	0.993	0.993		
Copper	N/A	N/A	0.83	0.83		
Lead	1.46203-[(In hardness)(0.145712)]	1.46203-[(In hardness)(0.145712)]	0.951	0.951		
Nickel	0.998	0.997	0.990	0.990		
Selenium	0.996	0.922	0.998	0.998		
Silver	0.85	0.85	0.85			
Zinc	0.978	0.986	0.946	0.946		

NOTE: Cadmium values  $m_A$ ,  $b_A$ , and the freshwater acute CF in Endnote F are not applicable until the freshwater acute cadmium criterion in Table 30 is approved by EPA and, if necessary, EPA withdraws the federal criteria for Oregon. Until then the federally promulgated freshwater acute cadmium criterion at 40 CFR 131.46 remains applicable for CWA purposes.

### **Endnote N: Deriving freshwater copper criteria**

The freshwater copper criteria at any time are the Biotic Ligand Model (BLM) derived Instantaneous Water Quality Criteria (IWQC) output based on a concurrently measured set of model input parameter values. The Biotic Ligand Model uses multiple ambient water quality parameters to derive 1-hour acute exposure (CMC) and 96-hour chronic exposure (CCC) water quality criteria (IWQC) for copper based on the site specific water chemistry that determines the toxicity of copper to aquatic life. If measured data for one or more of the model input parameters used to derive the acute and chronic IWQC is not available, the procedures in section (1) or (2) of this endnote will be used as specified to substitute an estimate or a default value for the missing input parameter. BLM results (IWQC) based on sufficient measured input parameter data are more accurate and supersede results based on estimates or default values. The acceptable BLM software to calculate the IWQC include version 2.2.3, referenced in "Aquatic Life Ambient Freshwater Quality Criteria – Copper": EPA-822-R-07-001, February 2007, and version 2.2.4. The criteria are expressed as dissolved copper in micrograms per liter (to the nearest one-tenth).

(1) Input Parameter Substitution and Estimation Procedures to Derive BLM Criteria (IWQC)

If the measured value for any input parameter needed to derive an IWQC using the BLM is not available, DEQ will substitute an estimated input parameter value according to the procedures described in this section [Endnote N (1)]. If the data required to determine the estimated parameter value is not available, DEQ will use default values derived according to the procedures in Endnote N (2).

- (a) Total recoverable concentration measurements will be substituted for dissolved concentration measurements that are not available. For alkalinity, calcium, chloride, magnesium, potassium, sodium and sulfate, total recoverable concentration measurements will be used as a direct substitute for dissolved concentration measurements. Total organic carbon (TOC) measurements will be multiplied by 0.83 to convert the TOC value to an equivalent dissolved organic carbon (DOC) value; except where sufficient TOC and DOC data are available for a site, DEQ will calculate and apply a site-specific translator in place of 0.83 to convert TOC values to DOC for use in the BLM.
- (b) Alkalinity, calcium, chloride, magnesium, potassium, sodium and sulfate: If data for any of these BLM input parameters are missing from a particular dataset, DEQ will estimate its value based on the relationship of the ion or alkalinity to specific conductance measurements for that data set using the regression analysis equations in Table 1. Specific conductance measurements must be concurrent with the other BLM input parameters dataset.

	Table N-1									
Parameter	Regression Equation									
Alkalinity	Alk. = $\exp^{(0.88 \cdot [\ln(SpC)] - 0.41)}$									
Calcium	Ca = $\exp^{(0.96 \cdot [\ln(SpC)] - 2.29)}$									
Chloride	$CI = exp^{(1.15\cdot[ln(SpC)] - 3.82)}$									
Magnesium	$Mg = exp^{(0.91 \cdot [ln(SpC)] - 3.09)}$									
Potassium	$K = exp^{(0.84 \cdot [ln(SpC)] - 3.74)}$									
Sodium	Na = $\exp^{(0.86 \cdot [\ln(SpC)] - 2.22)}$									
Sulfate	$SO_4 = exp^{(1.45 \cdot [ln(SpC)] - 5.59}$									

Where, "SpC" is a measurement of specific conductance in µmhos/cm, "In" is the natural logarithm, and "exp" is a mathematical constant that is the base of the natural logarithm.

### (c) pH

If concurrent pH data is missing from the sample dataset, DEQ will use a representative pH value determined by interpolating from data available for the site or proximate monitoring locations where conditions (such as type of water body, stream flow and geology) are similar to the site. DEQ will use the available data and methods to produce the best practicable estimate of pH for the site and time for which the IWQC is being derived.

### (d) Temperature

If concurrent temperature data is missing from the sample dataset, DEQ will use a monthly mean temperature based on data available for the site or proximate monitoring locations where conditions (such as type of water body and stream flow) are similar to the site.

### (e) Humic Acid

If sufficient high quality data on the percentage of humic acid as a proportion of DOC is available for a site, DEQ will use that value in the BLM in place of the default value of 10% used in the model.

### (2) Default Action Values

If the measured value for DOC, alkalinity, calcium, chloride, magnesium, potassium, sodium or sulfate is not available to derive an IWQC using the BLM, and the parameter value cannot be estimated as specified in section (1) above, DEQ will use a conservative input value for the missing parameter as described in this section [Endnote N (2)] to derive a default action value using the Biotic Ligand Model. The default action value will be used for Clean Water Act purposes until measured or estimated input parameter data are available to derive accurate copper criteria (IWQC) based on site specific water chemistry.

(a) The default input parameter values for DOC, alkalinity calcium, chloride, magnesium, potassium, sodium and sulfate will be the percentile value from the

distribution of the high quality data available for surface waters in the region as shown in Table N-2.

	Table N-2 Percentile of data distribution to be used as default value by region											
Region  DOC percentile  Alkalinity and lons percentile												
Willamette	20 <sup>th</sup>	20 <sup>th</sup>										
Coastal	20 <sup>th</sup>	20 <sup>th</sup>										
Cascades	20 <sup>th</sup>	20 <sup>th</sup>										
Eastern	15 <sup>th</sup>	15 <sup>th</sup>										
Columbia River	20 <sup>th</sup>	20 <sup>th</sup>										

- (b) The regional default values for each parameter and region will be updated periodically as additional high quality data becomes available and is added to DEQ's database.
- (c) The regional default values for each parameter are available on DEQ's website.
- (d) The regions listed in Table N-2 are comprised of the following EPA Level III ecoregions or waterbody:
  - (i) Willamette: the Willamette Valley
  - (ii) Coastal: Coast Range and Klamath Mountains
  - (iii) Cascades: Cascades
  - (iv) Eastern: Eastern Cascades Slopes and Foothills, Columbia Plateau, Blue

Mountains, Northern Basin and Range and Snake River Plain

(v) Columbia River: Columbia River mainstem in Oregon

### (3) General Policies

- (a) The copper BLM derives instantaneous criteria results (IWQC) that vary at a site over time reflecting the effect of local water chemistry on copper toxicity to aquatic organisms. DEQ will apply the BLM criteria for Clean Water Act purposes to protect the water body during the most bioavailable or toxic conditions.
- (b) For assessing waters of the state, DEQ will use approaches that give preference to the use of BLM criteria derived with site-specific measured input parameter data.

### **Endnote O: Deriving freshwater aluminum criteria**

Freshwater aluminum criteria values are calculated from a concurrently measured set of Aluminum Criteria Calculator (ACC) input parameter values. The input parameter values are based on the site and time specific water chemistry that determines the toxicity of aluminum to aquatic life. If measured data for one or more of the ACC input parameters is not available, the procedures in section (1), (2), or (3) of this endnote will be used, in the order listed, as specified to substitute an estimated or a default value for the missing input parameter or to apply default criteria derived using ecoregional data.

Criteria values based on sufficient concurrent measured input parameter data are more accurate, preferred, and supersede results based on estimates, default values or applied default ecoregional criteria values. The criteria are expressed as total recoverable in micrograms per liter (to two significant figures). Oregon will use analytical methods that measure the bioavailable fraction of aluminum unless total recoverable aluminum measurements are required by Federal regulations.

### (1) Input Parameter Estimation Procedures to Derive Aluminum Criteria Values

If the measured value for one or more input parameters needed to derive an aluminum criteria value using the ACC is not available, DEQ will substitute a calculated or estimated input value according to the procedures described in this section [Endnote O (1)].

### (a) DOC

DEQ will use total organic carbon (TOC) measurements to estimate DOC measurements that are not available. Total organic carbon (TOC) measurements will be multiplied by 0.83 to convert the TOC value to an equivalent dissolved organic carbon (DOC) value; except where sufficient TOC and DOC data are available for a site, DEQ will calculate and apply a site-specific translator in place of 0.83 to convert TOC values to DOC for use in the Aluminum Criteria Calculator. If neither DOC nor TOC measurements are available, substitute a default DOC value as described in Endnote O (2).

### (b) Total Hardness

If total hardness is not available, DEQ will estimate total hardness by substituting dissolved hardness as an input parameter for the Aluminum Criteria Calculator. If neither total nor dissolved hardness data are available, DEQ will use the equation in Table O-1 to estimate total hardness using specific conductance. Specific conductance measurements must be concurrent with the other input parameters for the Aluminum Criteria Calculator. If total hardness cannot be estimated from concurrent data, DEQ will apply the applicable ecoregional default aluminum criterion described in Endnote O (3).

Table O-1										
Equation to estimate total hardness from										
specific conductance										
Parameter Regression Equation										
Total Hardness = exp <sup>(1.050·[in(SpC)] - 1.211)</sup>										

Where, "SpC" is a measurement of specific conductance in µmhos/cm, "In" is the natural logarithm, and "exp" is a mathematical constant that is the base of the natural logarithm.

### (2) Applying a Default Value for DOC to Derive Aluminum Criteria Values

If concurrently measured DOC is not available to derive a criteria value using the ACC and DOC cannot be estimated as specified in Endnote O (1)(a) above, DEQ will use a conservative default DOC input value as described in this section [Endnote O (2)] to derive the criteria value.

The default DOC input value will be used for Clean Water Act purposes until measured or estimated DOC input data are available to derive aluminum criteria values based on site-specific water chemistry.

(a) The default input parameter values for DOC will be the percentile value from the distribution of the high-quality data available for surface waters in the region as shown in Table O-2.

Table O-2  Percentile of data distribution to be used as default value by region											
Region DOC percentile											
Willamette	15 <sup>th</sup>										
Coastal	30 <sup>th</sup>										
Cascades	20 <sup>th</sup>										
Eastern	15 <sup>th</sup>										
Columbia River	10 <sup>th</sup>										

- b) The regional default DOC values will be updated periodically as additional high-quality data become available and are added to DEQ's database.
- (c) The resulting regional default input values for DOC are shown on DEQ's website.
- (d) The regions listed in Table O-2 are the same as those defined in Endnote N(2)(d).

### (3) Applying Aluminum Default Ecoregional Criteria

If data for pH is missing or hardness is missing and cannot be estimated as described in Endnote O (1)(b), DEQ will apply an ecoregional default aluminum criteria value.

- (a) The default ecoregional acute (CMC) and chronic (CCC) criteria values will be the 10<sup>th</sup> percentile value from the distribution of all ACC outputs calculated from concurrently measured high quality input data available for Oregon surface waters by EPA Level III ecoregion with the Columbia River mainstem treated separately.
- (b) The ecoregional default aluminum criteria values will be updated periodically as additional high quality data become available and are added to DEQ's database.
- (c) The resulting ecoregional default aluminum criteria values are shown on DEQ's website.

### (4) General Policies

(a) The ACC produces criteria values that vary at a site over time reflecting the effect of local water chemistry on aluminum toxicity to aquatic organisms. To apply the aluminum

criteria for Clean Water Act purposes, criteria values based on ambient water chemistry conditions must protect the water body over the full range of water chemistry conditions, including during conditions when aluminum is most toxic.

(b) When applying the aluminum criteria, DEQ will use approaches that give preference to the use of ACC criteria values based on concurrently measured or estimated input parameter data, as described in Endnote O (1), (2), and (3) in the order listed, and concurrently measured aluminum data.

### **Table 30(a): Ammonia Acute Criteria Values (One-hour Average)**—Salmonid Species Present Temperature and pH-Dependent and expressed as Total Ammonia Nitrogen (mg/L TAN)

### Criteria cannot be exceeded more than once every three years

$$Acute \ Criterion = MIN \left( \left( \frac{0.275}{1 + 10^{7.204 - pH}} + \frac{39.0}{1 + 10^{pH - 7.204}} \right), \left( 0.7249 \times \left( \frac{0.0114}{1 + 10^{7.204 - pH}} + \frac{1.6181}{1 + 10^{pH - 7.204}} \right) \times \left( 23.12 \times 10^{0.036 \times (20 - T)} \right) \right) \right)$$

	Temperature (°C)																/ /
рН	0-14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
6.5	33	33	32	29	27	25	23	21	19	18	16	15	14	13	12	11	9.9
6.6	31	31	30	28	26	24	22	20	18	17	16	14	13	12	11	10	9.5
6.7	30	30	29	27	24	22	21	19	18	16	15	14	13	12	11	9.8	9.0
6.8	28	28	27	25	23	21	20	18	17	15	14	13	12	11	10	9.2	8.5
6.9	26	26	25	23	21	20	18	17	15	14	13	12	11	10	9.4	8.6	7.9
7.0	24	24	23	21	20	18	17	15	14	13	12	11	10	9.4	8.6	8.0	7.3
7.1	22	22	21	20	18	17	15	14	13	12	11	10	9.3	8.5	7.9	7.2	6.7
7.2	20	20	19	18	16	15	14	13	12	11	9.8	9.1	8.3	7.7	7.1	6.5	6.0
7.3	18	18	17	16	14	13	12	11	10	9.5	8.7	8.0	7.4	6.8	6.3	5.8	5.3
7.4	15	15	15	14	13	12	11	9.8	9.0	8.3	7.7	7.0	6.5	6.0	5.5	5.1	4.7
7.5	13	13	13	12	11	10	9.2	8.5	7.8	7.2	6.6	6.1	5.6	5.2	4.8	4.4	4.0
7.6	11	11	11	10	9.3	8.6	7.9	7.3	6.7	6.2	5.7	5.2	4.8	4.4	4.1	3.8	3.5
7.7	9.6	9.6	9.3	8.6	7.9	7.3	6.7	6.2	5.7	5.2	4.8	4.4	4.1	3.8	3.5	3.2	3.0
7.8	8.1	8.1	7.9	7.2	6.7	6.1	5.6	5.2	4.8	4.4	4.0	3.7	3.4	3.2	2.9	2.7	2.5
7.9	6.8	6.8	6.6	6.0	5.6	5.1	4.7	4.3	4.0	3.7	3.4	3.1	2.9	2.6	2.4	2.2	2.1
8.0	5.6	5.6	5.4	5.0	4.6	4.2	3.9	3.6	3.3	3.0	2.8	2.6	2.4	2.2	2.0	1.9	1.7
8.1	4.6	4.6	4.5	4.1	3.8	3.5	3.2	3.0	2.7	2.5	2.3	2.1	2.0	1.8	1.7	1.5	1.4
8.2	3.8	3.8	3.7	3.5	3.1	2.9	2.7	2.4	2.3	2.1	1.9	1.8	1.6	1.5	1.4	1.3	1.2
8.3	3.1	3.1	3.1	2.8	2.6	2.4	2.2	2.0	1.9	1.7	1.6	1.4	1.3	1.2	1.1	1.0	0.96
8.4	2.6	2.6	2.5	2.3	2.1	2.0	1.8	1.7	1.5	1.4	1.3	1.2	1.1	1.0	0.93	0.86	0.79
8.5	2.1	2.1	2.1	1.9	1.8	1.6	1.5	1.4	1.3	1.2	1.1	0.98	0.90	0.83	0.77	0.71	0.65
8.6	1.8	1.8	1.7	1.6	1.5	1.3	1.2	1.1	1.0	0.96	0.88	0.81	0.75	0.69	0.63	0.59	0.54
8.7	1.5	1.5	1.4	1.3	1.2	1.1	1.0	0.94	0.87	0.80	0.74	0.68	0.62	0.57	0.53	0.49	0.45
8.8	1.2	1.2	1.2	1.1	1.0	0.93	0.86	0.79	0.73	0.67	0.62	0.57	0.52	0.48	0.44	0.41	0.37
8.9	1.0	1.0	1.0	0.93	0.85	0.79	0.72	0.67	0.61	0.56	0.52	0.48	0.44	0.40	0.37	0.34	0.32
9.0	0.88	0.88	0.86	0.79	0.73	0.67	0.62	0.57	0.52	0.48	0.44	0.41	0.37	0.34	0.32	0.29	0.27

### Table 30(b): Ammonia Acute Criteria Values (One-hour Average\*)—Salmonid Species Absent

Temperature and pH-Dependent and expressed as Total Ammonia Nitrogen (mg/L TAN)

### Criteria cannot be exceeded more than once every three years

Acute Criterion =  $0.7249 \times \frac{0.0114}{1 + 10^{7.204 - pH}} + \frac{1.6181}{1 + 10^{pH - 7.204}} \times MIN(51.93, 23.12 \times 10^{0.036 \times (20 - T)})$ Temperature (°C)

												<b>O</b> )									
рН	0-10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
6.5	51	48	44	41	37	34	32	29	27	25	23	21	19	18	16	15	14	13	12	11	9.9
6.6	49	46	42	39	36	33	30	28	26	24	22	20	18	17	16	14	13	12	11	10	9.5
6.7	46	44	40	37	34	31	29	27	24	22	21	19	18	16	15	14	13	12	11	9.8	9.0
6.8	44	41	38	35	32	30	27	25	23	21	20	18	17	15	14	13	12	11	10	9.2	8.5
6.9	41	38	35	32	30	28	25	23	21	20	18	17	15	14	13	12	11	10	9.4	8.6	7.9
7.0	38	35	33	30	28	25	23	21	20	18	17	15	14	13	12	11	10	9.4	8.6	7.9	7.3
7.1	34	32	30	27	25	23	21	20	18	17	15	14	13	12	11	10	9.3	8.5	7.9	7.2	6.7
7.2	31	29	27	25	23	21	19	18	16	15	14	13	12	11	9.8	9.1	8.3	7.7	7.1	6.5	6.0
7.3	27	26	24	22	20	18	17	16	14	13	12	11	10	9.5	8.7	8.0	7.4	6.8	6.3	5.8	5.3
7.4	24	22	21	19	18	16	15	14	13	12	11	9.8	9.0	8.3	7.7	7.0	6.5	6.0	5.5	5.1	4.7
7.5	21	19	18	17	15	14	13	12	11	10	9.2	8.5	7.8	7.2	6.6	6.1	5.6	5.2	4.8	4.4	4.0
7.6	18	17	15	14	13	12	11	10	9.3	8.6	7.9	7.3	6.7	6.2	5.7	5.2	4.8	4.4	4.1	3.8	3.5
7.7	15	14	13	12	11	10	9.3	8.6	7.9	7.3	6.7	6.2	5.7	5.2	4.8	4.4	4.1	3.8	3.5	3.2	2.9
7.8	13	12	11	10	9.3	8.5	7.9	7.2	6.7	6.1	5.6	5.2	4.8	4.4	4.0	3.7	3.4	3.2	2.9	2.7	2.5
7.9	11	9.9	9.1	8.4	7.7	7.1	6.6	3.0	5.6	5.1	4.7	4.3	4.0	3.7	3.4	3.1	2.9	2.6	2.4	2.2	2.1
8.0	8.8	8.2	7.6	7.0	6.4	5.9	5.4	5.0	4.6	4.2	3.9	3.6	3.3	3.0	2.8	2.6	2.4	2.2	2.0	1.9	1.7
8.1	7.2	6.8	6.3	5.8	5.3	4.9	4.5	4.1	3.8	3.5	3.2	3.0	2.7	2.5	2.3	2.1	2.0	1.8	1.7	1.5	1.4
8.2	6.0	5.6	5.2	4.8	4.4	4.0	3.7	3.4	3.1	2.9	2.7	2.4	2.3	2.1	1.9	1.8	1.6	1.5	1.4	1.3	1.2
8.3	4.9	4.6	4.3	3.9	3.6	3.3	3.1	2.8	2.6	2.4	2.2	2.0	1.9	1.7	1.6	1.4	1.3	1.2	1.1	1.0	0.96
8.4	4.1	3.8	3.5	3.2	3.0	2.7	2.5	2.3	2.1	2.0	1.8	1.7	1.5	1.4	1.3	1.2	1.1	1.0	0.93	0.86	0.79
8.5	3.3	3.1	2.9	2.7	2.4	2.3	2.1	1.9	1.8	1.6	1.5	1.4	1.3	1.2	1.1	0.98	0.90	0.83	0.77	0.71	0.65
8.6	2.8	2.6	2.4	2.2	2.0	1.9	1.7	1.6	1.5	1.3	1.2	1.1	1.0	0.96	0.88	0.81	0.75	0.69	0.63	0.58	0.54
8.7	2.3	2.2	2.0	1.8	1.7	1.6	1.4	1.3	1.2	1.1	1.0	0.94	0.87	0.80	0.74	0.68	0.62	0.57	0.53	0.49	0.45
8.8	1.9	1.8	1.7	1.5	1.4	1.3	1.2	1.1	1.0	0.93	0.86	0.79	0.73	0.67	0.62	0.57	0.52	0.48	0.44	0.41	0.37
8.9	1.6	1.5	1.4	1.3	1.2	1.1	1.0	0.93	0.85	0.79	0.72	0.67	0.61	0.56	0.52	0.48	0.44	0.40	0.37	0.34	0.32
9.0	1.4	1.3	1.2	1.1	1.0	0.93	0.86	0.79	0.73	0.67	0.62	0.57	0.52	0.48	0.44	0.41	0.37	0.34	0.32	0.29	0.27

### Table 30(c): Ammonia Chronic Criteria Values (30-day Rolling Average\*)

Temperature and pH-Dependent and expressed as Total Ammonia Nitrogen (mg/L TAN)

\* The highest four-day average within the 30-day averaging period must not be more than 2.5 times the chronic value

### Criteria cannot be exceeded more than once every three years

Chronic Criterion = 
$$0.8876 \times \left(\frac{0.0278}{1 + 10^{7.688 - pH}} + \frac{1.1994}{1 + 10^{pH - 7.688}}\right) \times \left(2.126 \times 10^{0.028 \times \left(20 - MAX(T,7)\right)}\right)$$

									\ <u>1</u>	+ 10'		eratur		7.000	,									
рН	0-7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
6.5	4.9	4.6	4.3	4.1	3.8	3.6	3.3	3.1	2.9	2.8	2.6	2.4	2.3	2.1	2.0	1.9	1.8	1.6	1.5	1.5	1.4	1.3	1.2	1.1
6.6	4.8	4.5	4.3	4.0	3.8	3.5	3.3	3.1	2.9	2.7	2.5	2.4	2.2	2.1	2.0	1.8	1.7	1.6	1.5	1.4	1.3	1.3	1.2	1.1
6.7	4.8	4.5	4.2	3.9	3.7	3.5	3.2	3.0	2.8	2.7	2.5	2.3	2.2	2.1	1.9	1.8	1.7	1.6	1.5	1.4	1.3	1.2	1.2	1.1
6.8	4.6	4.4	4.1	3.8	3.6	3.4	3.2	3.0	2.8	2.6	2.4	2.3	2.1	2.0	1.9	1.8	1.7	1.6	1.5	1.4	1.3	1.2	1.1	1.1
6.9	4.5	4.2	4.0	3.7	3.5	3.3	3.1	2.9	2.7	2.5	2.4	2.2	2.1	2.0	1.8	1.7	1.6	1.5	1.4	1.3	1.2	1.2	1.1	1.0
7.0	4.4	4.1	3.8	3.6	3.4	3.2	3.0	2.8	2.6	2.4	2.3	2.2	2.0	1.9	1.8	1.7	1.6	1.5	1.4	1.3	1.2	1.1	1.1	0.99
7.1	4.2	3.9	3.7	3.5	3.2	3.0	2.8	2.7	2.5	2.3	2.2	2.1	1.9	1.8	1.7	1.6	1.5	1.4	1.3	1.2	1.2	1.1	1.0	0.95
7.2	4.0	3.7	3.5	3.3	3.1	2.9	2.7	2.5	2.4	2.2	2.1	2.0	1.8	1.7	1.6	1.5	1.4	1.3	1.3	1.2	1.1	1.0	0.96	0.90
7.3	3.8	3.5	3.3	3.1	2.9	2.7	2.6	2.4	2.2	2.1	2.0	1.8	1.7	1.6	1.5	1.4	1.3	1.3	1.2	1.1	1.0	0.97	0.91	0.85
7.4	3.5	3.3	3.1	2.9	2.7	2.5	2.4	2.2	2.1	2.0	1.8	1.7	1.6	1.5	1.4	1.3	1.3	1.2	1.1	1.0	0.96	0.90	0.85	0.79
7.5	3.2	3.0	2.8	2.7	2.5	2.3	2.2	2.1	1.9	1.8	1.7	1.6	1.5	1.4	1.3	1.2	1.2	1.1	1.0	0.95	0.89	0.83	0.78	0.73
7.6	2.9	2.8	2.6	2.4	2.3	2.1	2.0	1.9	1.8	1.6	1.5	1.4	1.4	1.3	1.2	1.1	1.1	0.98	0.92	0.86	0.81	0.76	0.71	0.67
7.7	2.6	2.4	2.3	2.2	2.0	1.9	1.8	1.7	1.6	1.5	1.4	1.3	1.2	1.1	1.1	1.0	0.94	0.88	0.83	0.78	0.73	0.68	0.64	0.60
7.8	2.3	2.2	2.1	1.9	1.8	1.7	1.6	1.5	1.4	1.3	1.2	1.2	1.1	1.0	0.95	0.89	0.84	0.79	0.74	0.69	0.65	0.61	0.57	0.53
7.9	2.1	1.9	1.8	1.7	1.6	1.5	1.4	1.3	1.2	1.2	1.1	1.0	0.95	0.89	0.84	0.79	0.74	0.69	0.65	0.61	0.57	0.53	0.50	0.47
8.0	1.8	1.7	1.6	1.5	1.4	1.3	1.2	1.1	1.1	1.0	0.94	0.88	0.83	0.78	0.73	0.68	0.64	0.60	0.56	0.53	0.50	0.44	0.44	0.41
8.1	1.5	1.5	1.4	1.3	1.2	1.1	1.1	0.99	0.92	0.87	0.81	0.76	0.71	0.67	0.63	0.59	0.55	0.52	0.49	0.46	0.43	0.40	0.38	0.35
8.2	1.3	1.2	1.2	1.1	1.0	0.96	0.90	0.84	0.79	0.74	0.70	0.65	0.61	0.57	0.54	0.50	0.47	0.44	0.42	0.39	0.37	0.34	0.32	0.30
8.3	1.1	1.1	0.99	0.93	0.87	0.82	0.76	0.72	0.67	0.63	0.59	0.55	0.52	0.49	0.46	0.43	0.40	0.38	0.35	0.33	0.31	0.29	0.27	0.26
8.4	0.95	0.89	0.84	0.79	0.74	0.69	0.65	0.61	0.57	0.53	0.50	0.47	0.44	0.41	0.39	0.36	0.34	0.32	0.30	0.28	0.26	0.25	0.23	0.22
8.5	0.80	0.75	0.71	0.67	0.62	0.58	0.55	0.51	0.48	0.45	0.42	0.40	0.37	0.35	0.33	0.31	0.29	0.27	0.25	0.24	0.22	0.21	0.20	0.18
8.6	0.68	0.64	0.60	0.56	0.53	0.49	0.46	0.43	0.41	0.38	0.36	0.33	0.31	0.29	0.28	0.26	0.24	0.23	0.21	0.20	0.19	0.18	0.16	0.15
8.7	0.57	0.54	0.51	0.47	0.44	0.42	0.39	0.37	0.34	0.32	0.30	0.28	0.27	0.25	0.23	0.22	0.21	0.19	0.18	0.17	0.16	0.15	0.14	0.13
8.8	0.49	0.46	0.43	0.40	0.38	0.35	0.33	0.31	0.29	0.27	0.26	0.24	0.23	0.21	0.20	0.19	0.17	0.16	0.15	0.14	0.13	0.13	0.12	0.11
8.9	0.42	0.39	0.37	0.34	0.32	0.30	0.28	0.27	0.25	0.23	0.22	0.21	0.19	0.18	0.17	0.16	0.15	0.14	0.13	0.12	0.12	0.11	0.10	0.09
9.0	0.36	0.34	0.32	0.30	0.28	0.26	0.24	0.23	0.21	0.20	0.19	0.18	0.17	0.16	0.15	0.14	0.13	0.12	0.11	0.11	0.10	0.09	0.09	0.08



Effective April 18, 2014

### **Human Health Criteria Summary**

The concentration for each pollutant listed in Table 40 was derived to protect Oregonians from potential adverse health impacts associated with long-term exposure to toxic substances associated with consumption of fish, shellfish, and water. The "organism only" criteria are established to protect fish and shellfish consumption and apply to waters of the state designated for fishing. The "water + organism" criteria are established to protect the consumption of drinking water, fish, and shellfish, and apply where both fishing and domestic water supply (public and private) are designated uses. All criteria are expressed as micrograms per liter (µg/L), unless otherwise noted. Pollutants are listed in alphabetical order. Additional information includes the Chemical Abstract Service (CAS) number, whether the criterion is based on carcinogenic effects (can cause cancer in humans), and whether there is an aquatic life criterion for the pollutant (i.e. "y"= yes, "n" = no). All the human health criteria were calculated using a fish consumption rate of 175 grams per day unless otherwise noted. A fish consumption rate of 175 grams per day is approximately equal to 23 8-ounce fish meals per month. For pollutants categorized as carcinogens, values represent a cancer risk of one additional case of cancer in one million people (i.e. 10<sup>-6</sup>), unless otherwise noted. All metals criteria are for total metal concentration, unless otherwise noted. Italicized pollutants represent non-priority pollutants.

					Human Health Criteria for Consumption of:						
No.	Pollutant	CAS Number	Carcinogen	Aquatic Life Criterion	Water + Organism (µg/L)	Organism Only (µg/L)					
1	Acenaphthene	83329	n	n	95	99					
2	Acrolein	107028	n	n	0.88	0.93					
3	Acrylonitrile	107131	у	n	0.018	0.025					
4	Aldrin	309002	у	у	0.0000050	0.0000050					
5	Anthracene	120127	n	n	2900	4000					
6	Antimony	7440360	n	n	5.1	64					
7	Arsenic (inorganic) <sup>A</sup>	7440382	у	у	2.1	2.1(freshwater) 1.0 (saltwater)					
	arsenic criteria are expresson of approximately 1 x 10 <sup>-5</sup> , an										
8	Asbestos <sup>B</sup>	1332214	у	n	7,000,000 fibers/L						
develo	human health risks from ast oped. The "water + organisr Orinking Water Act.										
9	Barium <sup>c</sup>	7440393	n	n	1000						
<sup>c</sup> The human health criterion for barium is the same as originally published in the 1976 EPA Red Book which predates the 1980 methodology and did not utilize the fish ingestion BCF approach. This same criterion value was also published in the 1986 EPA Gold Book. Human health risks are primarily from drinking water, therefore no "organism only" criterion was developed. The "water + organism" criterion is based on the Maximum Contaminant Level (MCL) established under the Safe Drinking Water Act.											
10	Benzene	71432	у	n	0.44	1.4					
11	Benzidine	92875	у	n	0.000018 0.0000						
	<del> </del>		i			i					

У

0.0013

0.0018

56553

12

Benz(a)anthracene

						Criteria for the ption of:
No.	Pollutant	CAS Number	Carcinogen	Aquatic Life Criterion	Water + Organism (µg/L)	Organism Only (µg/L)
13	Benzo(a)pyrene	50328	у	n	0.0013	0.0018
14	Benzo(b)fluoranthene 3,4	205992	у	n	0.0013	0.0018
15	Benzo(k)fluoranthene	207089	у	n	0.0013	0.0018
16	BHC Alpha	319846	у	n	0.00045	0.00049
17	BHC Beta	319857	у	n	0.0016	0.0017
18	BHC Gamma (Lindane)	58899	n	у	0.17	0.18
19	Bromoform	75252	у	n	3.3	14
20	Butylbenzyl Phthalate	85687	n	n	190	190
21	Carbon Tetrachloride	56235	у	n	0.10	0.16
22	Chlordane	57749	у	у	0.000081	0.000081
23	Chlorobenzene	108907	n	n	74	160
24	Chlorodibromomethan e	124481	у	n	0.31	1.3
25	Chloroethyl Ether bis 2	111444	у	n	0.020	0.053
26	Chloroform	67663	n	n	260	1100
27	Chloroisopropyl Ether bis 2	108601	n	n	1200	6500
28	Chloromethyl ether, bis	542881	у	n	0.000024	0.000029
29	Chloronaphthalene 2	91587	n	n	150	160

					Human Health Criteria for the Consumption of:			
No.	Pollutant	CAS Number	Carcinogen	Aquatic Life Criterion	Water + Organism (µg/L)	Organism Only (µg/L)		
30	Chlorophenol 2	95578	n	n	14	15		
31	Chlorophenoxy Herbicide (2,4,5,-TP) <sup>D</sup>	93721	n	n	10			

<sup>&</sup>lt;sup>D</sup> The Chlorophenoxy Herbicide (2,4,5,-TP) criterion is the same as originally published in the 1976 EPA Red Book which predates the 1980 methodology and did not utilize the fish ingestion BCF approach. This same criterion value was also published in the 1986 EPA Gold Book. Human health risks are primarily from drinking water, therefore no "organism only" criterion was developed. The "water + organism" criterion is based on the Maximum Contaminant Level (MCL) established under the Safe Drinking Water Act.

32	Chlorophenoxy Herbicide (2,4-D) <sup>E</sup>	94757	n	n	100	
----	---	-------	---	---	-----	--

<sup>&</sup>lt;sup>E</sup> The Chlorophenoxy Herbicide (2,4-D) criterion is the same as originally published in the 1976 EPA Red Book which predates the 1980 methodology and did not utilize the fish ingestion BCF approach. This same criterion value was also published in the 1986 EPA Gold Book. Human health risks are primarily from drinking water, therefore no "organism only" criterion was developed. The "water + organism" criterion is based on the Maximum Contaminant Level (MCL) established under the Safe Drinking Water Act.

33	Chrysene	218019	у	n	0.0013	0.0018
34	Copper <sup>F</sup>	7440508	n	у	1300	

F Human health risks from copper are primarily from drinking water, therefore no "organism only" criterion was developed. The "water + organism" criterion is based on the Maximum Contaminant Level (MCL) established under the Safe Drinking Water Act.

35	Cyanide <sup>G</sup>	57125	n	у	130	130		
	<sup>G</sup> The cyanide criterion is expressed as total cyanide (CN)/L.							
36	DDD 4,4'	72548	у	n	0.000031	0.000031		
37	DDE 4,4'	72559	у	n	0.000022	0.000022		
38	DDT 4,4'	50293	у	у	0.000022	0.000022		

					Human Health Criteria for the Consumption of:	
No.	Pollutant	CAS Number	Carcinogen	Aquatic Life Criterion	Water + Organism (µg/L)	Organism Only (µg/L)
39	Dibenz(a,h)anthracen e	53703	у	n	0.0013	0.0018
40	Dichlorobenzene(m) 1,3	541731	n	n	80	96
41	Dichlorobenzene(o) 1,2	95501	n	n	110	130
42	Dichlorobenzene(p) 1,4	106467	n	n	16	19
43	Dichlorobenzidine 3,3'	91941	у	n	0.0027	0.0028
44	Dichlorobromomethan e	75274	у	n	0.42	1.7
45	Dichloroethane 1,2	107062	у	n	0.35	3.7
46	Dichloroethylene 1,1	75354	n	n	230	710
47	Dichloroethylene trans 1,2	156605	n	n	120	1000
48	Dichlorophenol 2,4	120832	n	n	23	29
49	Dichloropropane 1,2	78875	у	n	0.38	1.5
50	Dichloropropene 1,3	542756	у	n	0.30	2.1
51	Dieldrin	60571	у	у	0.0000053	0.0000054
52	Diethyl Phthalate	84662	n	n	3800	4400
53	Dimethyl Phthalate	131113	n	n	84000	110000
54	Dimethylphenol 2,4	105679	n	n	76	85
55	Di-n-butyl Phthalate	84742	n	n	400	450

					Human Health Criteria for the Consumption of:	
No.	Pollutant	CAS Number	Carcinogen	Aquatic Life Criterion	Water + Organism (μg/L)	Organism Only (µg/L)
56	Dinitrophenol 2,4	51285	n	n	62	530
57	Dinitrophenols	2555058 7	n	n	62	530
58	Dinitrotoluene 2,4	121142	у	n	0.084	0.34
59	Dioxin (2,3,7,8-TCDD)	1746016	у	n	0.000000005 1	0.00000000051
60	Diphenylhydrazine 1,2	122667	у	n	0.014	0.020
61	Endosulfan Alpha	959988	n	у	8.5	8.9
62	Endosulfan Beta	3321365 9	n	у	8.5	8.9
63	Endosulfan Sulfate	1031078	n	n	8.5	8.9
64	Endrin	72208	n	у	0.024	0.024
65	Endrin Aldehyde	7421934	n	n	0.030	0.030
66	Ethylbenzene	100414	n	n	160	210
67	Ethylhexyl Phthalate bis 2	117817	у	n	0.20	0.22
68	Fluoranthene	206440	n	n	14	14
69	Fluorene	86737	n	n	390	530
70	Heptachlor	76448	у	у	0.0000079	0.0000079
71	Heptachlor Epoxide	1024573	у	у	0.0000039	0.0000039
72	Hexachlorobenzene	118741	у	n	0.000029	0.000029

					Human Health Criteria for the Consumption of:	
No.	Pollutant	CAS Number	Carcinogen	Aquatic Life Criterion	Water + Organism (µg/L)	Organism Only (μg/L)
73	Hexachlorobutadiene	87683	у	n	0.36	1.8
74	Hexachlorocyclo- hexane-Technical	608731	у	n	0.0014	0.0015
75	Hexachlorocyclopenta diene	77474	n	n	30	110
76	Hexachloroethane	67721	у	n	0.29	0.33
77	Indeno(1,2,3- cd)pyrene	193395	у	n	0.0013	0.0018
78	Isophorone	78591	у	n	27	96
79	Manganese H	7439965	n	n		100

<sup>&</sup>lt;sup>H</sup> The "fish consumption only" criterion for manganese applies only to salt water and is for total manganese. This EPA recommended criterion predates the 1980 human health methodology and does not utilize the fish ingestion BCF calculation method or a fish consumption rate.

80	Methoxychlor <sup>I</sup>	72435	n	у	100	
----	---------------------------	-------	---	---	-----	--

<sup>&</sup>lt;sup>1</sup> The human health criterion for methoxychlor is the same as originally published in the 1976 EPA Red Book which predates the 1980 methodology and did not utilize the fish ingestion BCF approach. This same criterion value was also published in the 1986 EPA Gold Book. Human health risks are primarily from drinking water, therefore no "organism only" criterion was developed. The "water + organism" criterion is based on the Maximum Contaminant Level (MCL) established under the Safe Drinking Water Act.

81	Methyl Bromide	74839	n	n	37	150
82	Methyl-4,6- dinitrophenol 2	534521	n	n	9.2	28
83	Methylene Chloride	75092	у	n	4.3	59
84	Methylmercury (mg/kg) <sup>J</sup>	2296792 6	n	n		0.040 mg/kg

					Human Health Criteria for the Consumption of:			
No.	Pollutant	CAS Number	Carcinogen	Aquatic Life Criterion	Water + Organism (µg/L)	Organism Only (μg/L)		
J This value is expressed as the fish tissue concentration of methylmercury. Contaminated fish and shellfish is the primary human route of exposure to methylmercury.								
85	Nickel	7440020	n	У	140	170		
96	Nitratos K	1479755	n	n	10000			

<sup>&</sup>lt;sup>K</sup> The human health criterion for nitrates is the same as originally published in the 1976 EPA Red Book which predates the 1980 methodology and did not utilize the fish ingestion BCF approach. This same criterion value was also published in the 1986 EPA Gold Book. Human health risks are primarily from drinking water, therefore no "organism only" criterion was developed. The "water + organism" criterion is based on the Maximum Contaminant Level (MCL) established under the Safe Drinking Water Act.

8

10000

86

Nitrates K

87	Nitrobenzene	98953	n	n	14	69
88	Nitrosamines	3557691 1	у	n	0.00079	0.046
89	Nitrosodibutylamine, N	924163	у	n	0.0050	0.022
90	Nitrosodiethylamine, N	55185	у	n	0.00079	0.046
91	Nitrosodimethylamine, N	62759	у	n	0.00068	0.30
92	Nitrosodi-n- propylamine, N	621647	у	n	0.0046	0.051
93	Nitrosodiphenylamine, N	86306	у	n	0.55	0.60
94	Nitrosopyrrolidine, N	930552	у	n	0.016	3.4
95	Pentachlorobenzene	608935	n	n	0.15	0.15
96	Pentachlorophenol	87865	у	у	0.15	0.30

					Human Health Criteria for the Consumption of:				
No.	Pollutant	CAS Number	Carcinogen	Aquatic Life Criterion	Water + Organism (µg/L)	Organism Only (µg/L)			
97	Phenol	108952	n	n	9400	86000			
98	Polychlorinated Biphenyls (PCBs) <sup>L</sup>	NA	у	у	0.0000064	0.0000064			
	<sup>L</sup> This criterion applies to total PCBs (e.g. determined as Aroclors or congeners).								
99	Pyrene	129000	n	n	290	400			
100	Selenium	7782492	n	у	120	420			
101	Tetrachlorobenzene, 1,2,4,5-	95943	n	n	0.11	0.11			
102	Tetrachloroethane 1,1,2,2	79345	у	n	0.12	0.40			
103	Tetrachloroethylene	127184	у	n	0.24	0.33			
104	Thallium	7440280	n	n	0.043	0.047			
105	Toluene	108883	n	n	720	1500			
106	Toxaphene	8001352	у	у	0.000028	0.000028			
107	Trichlorobenzene 1,2,4	120821	n	n	6.4	7.0			
108	Trichloroethane 1,1,2	79005	у	n	0.44	1.6			
109	Trichloroethylene	79016	у	n	1.4	3.0			
110	Trichlorophenol 2,4,6	88062	у	n	0.23	0.24			
111	Trichlorophenol, 2, 4, 5-	95954	n	n	330	360			
112	Vinyl Chloride	75014	у	n	0.023	0.24			

					Human Health Criteria for the Consumption of:	
No.	Pollutant	CAS Number	Carcinogen	Aquatic Life Criterion	Water + Organism (μg/L)	Organism Only (μg/L)
113	Zinc	7440666	n	у	2100	2600